# **Southeast Conservation Blueprint Summary**

for Florida

### Created 01/19/2024

## **Table of Contents**

About the Southeast Blueprint	<u> </u>
Southeast Blueprint Priorities	Δ
Hubs and Corridors	6
Indicator Summary	3
Threats	81
Ownership and Partners	85
Credits	90

The Southeast Conservation Adaptation Strategy

**SECAS** 



The Southeast Conservation Blueprint 2023

Southeast Conservation Blueprint Summary for Florida				
[	[THIS PAGE INTENTIONALLY LEFT BLANK]			

## **About the Southeast Blueprint**

The Southeast Conservation Blueprint is the primary product of the <u>Southeast Conservation Adaptation Strategy</u> (SECAS). It is a living, spatial plan to achieve the SECAS vision of a connected network of lands and waters across the Southeast and Caribbean. The Blueprint is regularly updated to incorporate new data, partner input, and information about on-the-ground conditions.

The Blueprint identifies priority areas based on a suite of natural and cultural resource indicators representing terrestrial, freshwater, and marine ecosystems. A connectivity analysis identifies corridors that link coastal and inland areas and span climate gradients.

### For more information:

- Visit the <u>Blueprint webpage</u>
- Review the <u>Blueprint 2023 Development Process</u>
- View and download the Blueprint data and make maps on the Blueprint page of the SECAS Atlas


### We're here to help!

- Do you have a question about the Blueprint?
- Would you like help using the Blueprint to support a proposal or inform a decision?
- Do you have a suggestion on how to improve the Blueprint? The Blueprint and its inputs are regularly revised based on input from people like you.
- Do you have feedback on how to improve the Simple Viewer interface?

If you need help or have questions, <u>contact Southeast Blueprint staff</u> by reaching out to a member of the user support team.

We're here to support you. We really mean it. It's what we do!

# **Southeast Blueprint Priorities**



### **Priority Categories**

### For a connected network of lands and waters

In total, Blueprint priorities and priority connections cover roughly 50% of the Southeast Blueprint geography.

### **Highest priority**

Areas where conservation action would make the biggest impact, based on a suite of natural and cultural resource indicators. This class covers roughly 10% of the Southeast Blueprint geography.

### **High priority**

Areas where conservation action would make a big impact, based on a suite of natural and cultural resource indicators. This class covers roughly 15% of the Southeast Blueprint geography.

### **Medium priority**

Areas where conservation action would make an above-average impact, based on a suite of natural and cultural resource indicators. This class covers roughly 20% of the Southeast Blueprint geography.

### **Priority connections**

Connections between priority areas that cover the shortest distance possible while routing through as much Blueprint priority as possible. This class covers roughly 5% of the Southeast Blueprint geography.

Table 1: Extent of each Blueprint priority category within Florida.

Priority Category	Acres	Percent of Area
Highest priority	5,061,404	11.1%
High priority	9,202,761	20.1%
Medium priority	10,583,796	23.2%
Priority connections	2,467,578	5.4%
Lower priority	18,382,624	40.2%
Total area	45,698,163	100%

### **Hubs and Corridors**

The Blueprint uses a least-cost path connectivity analysis to identify corridors that link hubs across the shortest distance possible, while also routing through as much Blueprint priority as possible.

Inland hubs are large patches (~5,000+ acres) of highest priority Blueprint areas and/or protected lands, connected by inland corridors. Marine and estuarine hubs are large estuaries and large patches (~5,000+ acres) of highest priority Blueprint areas. Marine and estuarine corridors connect those hubs within broad marine mammal movement areas.




Table 2: Extent of hubs and corridors within Florida.

Туре	Acres	Percent of Area
Inland continental hubs	10,894,379	23.8%
Inland continental corridors	4,215,242	9.2%
Marine & estuarine continental hubs	2,899,049	6.3%
Marine & estuarine continental corridors	3,369,573	7.4%
Not a hub or corridor	24,319,919	53.2%
Total area	45,698,163	100%

# **Indicator Summary**

Table 3: Terrestrial indicators.

Indicator	Present
East Coastal Plain open pine birds	✓
Equitable access to potential parks	<b>✓</b>
Fire frequency	<b>√</b>
Greenways & trails	<b>√</b>
<u>Intact habitat cores</u>	<b>√</b>
Interior Southeast grasslands	<b>✓</b>
Resilient terrestrial sites	<b>✓</b>
South Atlantic amphibian & reptile areas	<b>✓</b>
South Atlantic forest birds	<b>✓</b>
South Atlantic low-urban historic landscapes	✓
<u>Urban park size</u>	<b>√</b>

### Table 4: Freshwater indicators.

Indicator	Present
Atlantic migratory fish habitat	<b>✓</b>
Gulf migratory fish connectivity	<b>√</b>
Imperiled aquatic species	✓
Natural landcover in floodplains	✓
Network complexity	<b>✓</b>
Permeable surface	<b>√</b>

Table 5: Coastal & marine indicators.

Indicator	Present
Atlantic coral & hardbottom	✓
Atlantic deep-sea coral richness	✓
Atlantic estuarine fish habitat	✓
Atlantic marine birds	✓
Atlantic marine mammals	✓
<u>Coastal shoreline condition</u>	✓
Estuarine coastal condition	✓
Gulf coral & hardbottom	✓
Gulf deep-sea coral richness	✓
Gulf marine mammals	✓
Gulf sea turtles	✓
Island habitat	✓
Marine highly migratory fish	✓
Resilient coastal sites	✓
<u>Seagrass</u>	✓
South Atlantic beach birds	✓
South Atlantic maritime forest	✓
Stable coastal wetlands	✓



## East Coastal Plain open pine birds

This indicator identifies areas within the historic longleaf pine range east of the Mississippi River where creating or maintaining open pine habitat would most benefit six focal species of birds (Bachman's sparrow, red-cockaded woodpecker, Henslow's sparrow, red-headed woodpecker, Northern bobwhite, brown-headed nuthatch). It prioritizes areas for open pine conservation based on suitability for longleaf pine, feasibility of prescribed burning, proximity to protected lands, habitat suitability for focal bird species, and proximity to bird source populations. It originates from the East Gulf Coastal Plain Joint Venture's prioritization of areas for open pine ecosystem restoration.





Table 6: Indicator values for east coastal plain open pine birds within Florida. A good condition threshold is not yet defined for this indicator.

	Indicator Values: Priority for open pine conservation for focal bird species	Acres	Percent of Area
↑ High	High priority (score >80-100)	141,076	0.3%
	Medium-high priority (score >60-80)	401,386	0.9%
	Medium priority (score >40-60)	1,227,550	2.7%
	Medium-low priority (score >20-40)	3,593,119	7.9%
	Low priority (score 0-20)	564,675	1.2%
↓ Low	Not a priority (not identified as upland pine)	21,037,779	46.0%
	Area not evaluated for this indicator	18,732,576	41.0%
	Total area	45,698,163	100%



## **Equitable access to potential parks**

This cultural resource indicator prioritizes places to create new parks that would fill gaps in equitable access to open space within socially vulnerable communities in urban areas. It identifies areas where residents currently lack access to parks within a 10-minute walk (accounting for walkable road networks and access barriers like highways and fences), then prioritizes based on park need using demographic and environmental metrics. Parks help improve public health, foster a conservation ethic by providing opportunities for people to connect with nature, and support critical ecosystem services. This indicator originates from the Trust for Public Land's ParkServe park priority areas and the Center for Disease Control's Social Vulnerability Index.





High priority

Moderate priority

Not identified as a priority (within urban areas)

Table 7: Indicator values for equitable access to potential parks within Florida. A good condition threshold is not yet defined for this indicator.

	Indicator Values: Priority for a new park that would create nearby equitable access	Acres	Percent of Area
↑ High	Very high priority	242,084	0.5%
	High priority	251,217	0.5%
	Moderate priority	322,143	0.7%
↓ Low	Not identified as a priority (within urban areas)	36,486,525	79.8%
	Area not evaluated for this indicator	8,396,193	18.4%
	Total area	45,698,163	100%



This indicator uses remote sensing to estimate the number of times an area has been burned from 2013 to 2021. Many Southeastern ecosystems rely on regular, low-intensity fires to maintain habitat, encourage native plant growth, and reduce wildfire risk. This indicator combines burned area layers from U.S. Geological Survey Landsat data and the inter-agency Monitoring Trends in Burn Severity program. Landsat-based fire predictions within the range of longleaf pine are also available through <u>Southeast FireMap</u>.





Table 8: Indicator values for fire frequency within Florida. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.

	Indicator Values	Acres	Percent of Area	
↑ High	Burned 3+ times from 2013-2021	233,686	0.5%	
	Burned 2 times from 2013-2021	798,439	1.7%	↑ In good condition
	Burned 1 time from 2013-2021	2,400,843	5.3%	↓ Not in good
↓ Low	Not burned from 2013-2021 or row crop	38,608,848	84.5%	condition
	Area not evaluated for this indicator	3,656,346	8.0%	
	Total area	45,698,163	100%	



This cultural resource indicator measures both the natural condition and connected length of greenways and trails to characterize the quality of the recreational experience. Natural condition is based on the amount of impervious surface surrounding the path. Connected length captures how far a person can go without leaving a dedicated path, based on common distances for walking, running, and biking. This indicator originates from OpenStreetMap data and the National Land Cover Database.





Table 9: Indicator values for greenways & trails within Florida. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.

	Indicator Values	Acres	Percent of Area	
↑ High	Mostly natural and connected for ≥40 km	15,062	<0.1%	
	Mostly natural and connected for 5 to <40 km or partly natural and connected for ≥40 km	28,770	<0.1%	
	Mostly natural and connected for 1.9 to <5 km, partly natural and connected for 5 to <40 km, or developed and connected for ≥40 km	50,065	0.1%	
	Mostly natural and connected for <1.9 km, partly natural and connected for 1.9 to <5 km, or developed and connected for 5 to <40 km	23,292	<0.1%	↑ In good condition
	Partly natural and connected for <1.9 km or developed and connected for 1.9 to <5 km	12,620	<0.1%	→ Not in good condition
	Developed and connected for <1.9 km	19,977	<0.1%	
	Sidewalk	211,412	0.5%	
↓ Low	Not identified as a trail, sidewalk, or other path	41,570,672	91.0%	
	Area not evaluated for this indicator	3,766,292	8.2%	
	Total area	45,698,163	100%	



This indicator represents the size of large, unfragmented patches of natural habitat. It identifies minimally disturbed natural areas at least 100 acres in size and greater than 200 meters wide. Large areas of intact natural habitat are important for many wildlife species, including reptiles and amphibians, birds, and large mammals. This indicator originates from Esri's green infrastructure data.




Table 10: Indicator values for intact habitat cores within Florida. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.

	Indicator Values	Acres	Percent of Area	
↑ High	Large core (>10,000 acres)	7,882,144	17.2%	
	Medium core (>1,000-10,000 acres)	7,476,883	16.4%	
	Small core (>100-1,000 acres)	4,272,359	9.3%	↑ In good condition
↓ Low	Not a core	22,300,485	48.8%	→ Not in good condition
	Area not evaluated for this indicator	3,766,292	8.2%	
	Total area	45,698,163	100%	

# Terrestrial Intorio

## Interior Southeast grasslands

This indicator represents grasslands in the interior southeastern United States, which support important plants, birds, and pollinators. It includes grasslands with and without trees that are historically maintained by geology (e.g., outcrops, glades, and barrens), fire (e.g., Piedmont prairies), and/or the regular violent flooding on the banks of high-energy rivers known as "riverscour" (e.g., riverscour prairies). Known grasslands receive the highest scores, followed by bumble bee habitat buffers around known sites, areas in potentially compatible management, and restoration opportunities within grassland geology. This indicator combines data from multiple sources, including the Southeastern Grasslands Institute, Central Hardwoods Joint Venture, Rangeland Analysis Platform, and more.




Table 11: Indicator values for Interior Southeast grasslands within Florida. A good condition threshold is not yet defined for this indicator.

	Indicator Values	Acres	Percent of Area
↑ High	Known grassland	0	0%
	Known grassland buffer	0	0%
	Potentially compatible management within grassland geology (undeveloped powerline right-of-way or perennial forbs and grasses)	67	<0.1%
	Potentially compatible management outside of grassland geology (undeveloped powerline right-of-way or perennial forbs and grasses)	338,117	0.7%
	Grassland geology	316	<0.1%
↓ Low	Grassland less likely	4,161,358	9.1%
	Area not evaluated for this indicator	41,198,304	90.2%
	Total area	45,698,163	100%

# Terrestrial Resilient terrestrial sites

This indicator depicts an area's capacity to maintain species diversity and ecosystem function in the face of climate change. It measures two factors that influence resilience. The first, landscape diversity, reflects the number of microhabitats and climatic gradients created by topography, elevation, and hydrology. The second, local connectedness, reflects the degree of habitat fragmentation and strength of barriers to species movement. Highly resilient sites contain many different habitat niches that support biodiversity, and allow species to move freely through the landscape to find suitable microclimates as the climate changes. This indicator originates from The Nature Conservancy's Resilient Land data.




Table 12: Indicator values for resilient terrestrial sites within Florida. A good condition threshold is not yet defined for this indicator.

	Indicator Values	Acres	Percent of Area
↑ High	Most resilient	540,595	1.2%
	More resilient	4,110,579	9.0%
	Slightly more resilient	3,773,682	8.3%
	Average/median resilience	8,189,379	17.9%
	Slightly less resilient	3,179,043	7.0%
	Less resilient	3,497,260	7.7%
	Least resilient	866,993	1.9%
↓ Low	Developed	5,375,658	11.8%
	Area not evaluated for this indicator	16,164,974	35.4%
	Total area	45,698,163	100%

# Terrestrial

# South Atlantic amphibian & reptile areas

This indicator represents Priority Amphibian and Reptile Conservation Areas (PARCAs) in the South Atlantic. PARCA is an expert-driven, nonregulatory designation that includes places capable of supporting viable amphibian and reptile populations, places occupied by rare or imperiled species, and places rich in biodiversity or species unique to that geographic area (i.e., endemism). Reptiles and amphibians are a critical part of the Southeast region's rich biodiversity and many populations are declining in the face of threats like habitat loss, invasive species, and climate change.




Table 13: Indicator values for South Atlantic amphibian & reptile areas within Florida. A good condition threshold is not yet defined for this indicator.

	Indicator Values	Acres	Percent of Area
↑ High	Priority Amphibian and Reptile Conservation Area (PARCA)	2,908,341	6.4%
↓ Low	Not a Priority Amphibian and Reptile Conservation Area (PARCA)	7,359,596	16.1%
	Area not evaluated for this indicator	35,430,225	77.5%
	Total area	45,698,163	100%

### **Priority Amphibian and Reptile Conservation Areas:**

### **Apalachicola**

Extensive areas of longleaf pine savannas and sandhill habitat with a variety of wetlands, rivers, and streams; frosted flatwoods salamander (largest population with ca. 60 ponds), gopher frog, striped newt (once known from 20 ponds, but populations have seriously declined), gopher tortoise, Apalachicola dusky salamander, Barbour's map turtle, alligator snapping turtle, coal skink, Apalachicola kingsnake, southern copperhead.

### **Camp Blanding**

Extensive sandhill habitat with assorted wetlands; species of note include striped newt, gopher frog, gopher tortoise, Florida pine snake, eastern indigo snake.

### Florida Red Hills

Upland pine forest with gopher tortoises, primarily on quail plantations. It's not clear what is important about this area – no indigo records, one extinct striped newt pond, no gopher frog records; it is good for diamondback rattlesnakes, but they occur almost everywhere in Florida.

### **Georgia Barrier Islands and Marshes**

Nesting (island beaches) and/or foraging habitat (estuaries and nearshore waters) for four marine turtles (green, loggerhead, Kemp's ridley, and leatherback). Estuaries and embedded marsh islands are habitat for diamondback terrapins. Other rare species found in upland areas in this region include island glass lizards and dense populations of eastern diamondback rattlesnakes.

### **Lake Seminole Region**

Longleaf pine communities and embedded isolated wetlands provide habitat for gopher tortoises and eastern diamondback rattlesnakes. The Lower Chattahoochee and Flint Rivers, as well as Spring Creek, are inhabited by good populations of Barbour's map and alligator snapping turtles. Apalachicola dusky and Chamberlain's dwarf salamanders are found in seepages in this region. This area is underlain by the Floridan aquifer, which is home to the Georgia blind salamander.

### **Okefenokee Swamp**

This is the largest wetland in Georgia and includes both embedded (islands) and adjacent upland habitats. Striped crayfish snakes and Florida red-bellied turtles, found in very few other places in Southeast Georgia, thrive here. Frosted flatwoods salamanders, striped newts, gopher frogs, many-lined salamanders, dwarf sirens, carpenter Frogs, gopher tortoises, spotted turtles, eastern indigo snakes, eastern diamondback rattlesnakes, Florida green watersnakes, and perhaps mimic glass lizards all occur here.

#### St. Marks

Pine flatwoods, sandhills, and coastal habitats that support the second largest population of the frosted flatwoods salamander (>40 breeding ponds), but striped newts and gopher frogs were last recorded here in 1979. Species of note include the one-toed amphiuma, gopher tortoise, eastern kingsnake, Gulf salt marsh snake, and the westernmost record of the spotted turtle.


### Suwannee/Santa Fe/Osceola

Extensive pine flatwoods and swamps (Osceola) and sandhill habitat (along the Suwannee and Santa Fe Rivers). Species of note include the easternmost population of the alligator snapping turtle (may be a separate taxon), Suwannee cooter, indigo snake, tiger salamander, gopher tortoise, southern hognose snake, and Florida pine snake along the Suwannee and Santa Fe rivers, in addition to the many-lined salamander, carpenter frog, timber rattlesnake, spotted turtle, and possibly the last remaining population of the frosted flatwoods salamander on public land in the peninsula (Osceola).



### South Atlantic forest birds

This indicator is an index of habitat suitability for twelve upland hardwood and forested wetland bird species (wood thrush, whip-poor-will, American woodcock, red-headed woodpecker, Chuck-will's widow, hooded warbler, Kentucky warbler, Acadian flycatcher, Northern parula, black-throated green warbler, prothonotary warbler, Swainson's warbler) based on patch size and other ecosystem characteristics such as proximity to water and proximity to forest and ecotone edge. The needs of these species are increasingly restrictive at higher index values, reflecting better quality habitat. It originates from Southeast Gap Analysis Program and Designing Sustainable Landscapes bird habitat models.



warbler or Kentucky warbler)

or American woodcock)

Very small patches or near open areas (potential for presence of wood thrush, whip-poor-will, red-headed woodpecker, Chuck-will's widow,

Less potential for presence of forest bird index species

Table 14: Indicator values for South Atlantic forest birds within Florida. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.

	Indicator Values	Acres	Percent of Area	
↑ High	Very large patches near water (potential for presence of Swainson's warbler)	184,657	0.4%	
	Large patches often near water (potential for presence of Northern parula, black-throated green warbler, or prothonotary warbler)	1,629,936	3.6%	
	Medium patches (potential for presence of Acadian flycatcher)	431,559	0.9%	
	Small patches often near water (potential for presence of hooded warbler or Kentucky warbler)	103,594	0.2%	↑ In good condition
	Very small patches or near open areas (potential for presence of wood thrush, whippoor-will, red-headed woodpecker, Chuckwill's widow, or American woodcock)	5,789,774	12.7%	↓ Not in good condition
↓ Low	Less potential for presence of forest bird index species	2,044,572	4.5%	
	Area not evaluated for this indicator	35,514,072	77.7%	
	Total area	45,698,163	100%	

# Terrestrial

# South Atlantic low-urban historic landscapes

This cultural resource indicator is an index of sites on the National Register of Historic Places surrounded by limited urban development. It identifies significant historic places that remain connected to their context in the natural world. It uses the National Land Cover Database and historic places data from the National Park Service and various state historic resource agencies.



Table 15: Indicator values for South Atlantic low-urban historic landscapes within Florida. A good condition threshold is not yet defined for this indicator.

	Indicator Values	Acres	Percent of Area
↑ High	Historic place with nearby low-urban buffer	23,523	<0.1%
	Historic place with nearby high-urban buffer	2,915	<0.1%
↓ Low	Not in the National Register of Historic Places	9,965,648	21.8%
	Area not evaluated for this indicator	35,706,076	78.1%
	Total area	45,698,163	100%



This cultural resource indicator measures the size of parks larger than 5 acres in the urban environment. Protected natural areas in urban environments provide urban residents a nearby place to connect with nature, and offer refugia for some species. This indicator complements the equitable access to potential parks indicator by capturing the value of existing parks. It originates from the Protected Areas Database of the United States, Census urban areas, and the National Land Cover Database.




Table 16: Indicator values for urban park size within Florida. A good condition threshold is not yet defined for this indicator.

	Indicator Values	Acres	Percent of Area
↑ High	75+ acre urban park	1,084,548	2.4%
	50 to <75 acre urban park	19,379	<0.1%
	30 to <50 acre urban park	21,675	<0.1%
	10 to <30 acre urban park	33,605	<0.1%
	5 to <10 acre urban park	15,349	<0.1%
↓ Low	<5 acre urban park or not identified as an urban park	40,811,851	89.3%
	Area not evaluated for this indicator	3,711,754	8.1%
	Total area	45,698,163	100%

# Freshwater Atlantic migratory fish habitat

This indicator measures the condition of migratory fish habitat along the Atlantic coast within each catchment, using metrics of water quality, aquatic connectivity, habitat fragmentation, flow alteration, and more. Areas of excellent fish habitat are already in good condition and face few threats. Restoration opportunity areas are doing well in some respects, but restoration projects could significantly improve them. Degraded areas of opportunity face many challenges, and restoration projects are unlikely to increase available fish habitat unless particularly large in scope and scale. This indicator originates from the Atlantic Coast Fish Habitat Partnership's fish habitat conservation area mapping and prioritization project.




Table 17: Indicator values for Atlantic migratory fish habitat within Florida. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.

	Indicator Values	Acres	Percent of Area	
↑ High	Final score of 80 (areas of excellent fish habitat)	12,580	<0.1%	
	Final score of 70 (areas of excellent fish habitat)	5,591	<0.1%	↑ In good condition
	Final score of 60 (restoration opportunity areas)	905,271	2.0%	↓ Not in good condition
	Final score of 50 (restoration opportunity areas)	262,010	0.6%	
	Final score of 40 (restoration opportunity areas)	314,311	0.7%	
	Final score of 30 (restoration opportunity areas)	69,171	0.2%	
	Final score of 20 (restoration opportunity areas)	36,262	<0.1%	
	Final score of 10 (degraded areas of opportunity)	1,416	<0.1%	
↓ Low	Final score of 0 (degraded areas of opportunity)	0	0%	
	Area not evaluated for this indicator	44,091,550	96.5%	
	Total area	45,698,163	100%	

# Freshwater Gulf m

# **Gulf migratory fish connectivity**

This indicator captures how far upstream migratory fish in the Gulf of Mexico have been observed. How far upstream migratory fish can travel reflects not just the presence of dams and other barriers, but also the presence of measures like fish ladders that allow specific species to access habitat upstream of dams. This indicator originates from The Nature Conservancy's Southeast Aquatic Connectivity Assessment Project and applies to the Environmental Protection Agency's estimated floodplain, which spatially defines areas estimated to be inundated by a 100-year flood (also known as the 1% annual chance flood).




Table 18: Indicator values for Gulf migratory fish connectivity within Florida. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.

	Indicator Values	Acres	Percent of Area	
↑ High	Presence of Gulf sturgeon	1,353,893	3.0%	
	Presence of Alabama shad, American shad, or striped bass	132,387	0.3%	↑ In good condition
↓ Low	Not identified as Gulf migratory fish habitat (east of the Mississippi River)	12,091,417	26.5%	→ Not in good condition
	Area not evaluated for this indicator	32,120,465	70.3%	
	Total area	45,698,163	100%	

# Freshwater Imperiled aquatic species

This indicator measures the number of aquatic animal Species of Greatest Conservation Need (SGCN) observed within each 12-digit HUC subwatershed, including fish, mussels, snails, crayfish, and amphibians. SGCN are identified in State Wildlife Action Plans as most in need of conservation action. This indicator captures patterns of rare and endemic aquatic species diversity. It originates from state Natural Heritage Program data collected by the Southeast Aquatic Resources Partnership and applies to the Environmental Protection Agency's estimated floodplain, which spatially defines areas estimated to be inundated by a 100-year flood (also known as the 1% annual chance flood).







Table 19: Indicator values for imperiled aquatic species within Florida. A good condition threshold is not yet defined for this indicator.

	Indicator Values: Number of aquatic animal Species of Greatest Conservation Need observed	Acres	Percent of Area
↑ High	8+ species	51,356	0.1%
	7 species	34,114	<0.1%
	6 species	115,507	0.3%
	5 species	90,517	0.2%
	4 species	106,927	0.2%
	3 species	137,738	0.3%
	2 species	147,128	0.3%
	1 species	647,203	1.4%
	0 species	12,118,968	26.5%
↓ Low	Not identified as a floodplain (excluding West Virginia)	22,838,434	50.0%
	Area not evaluated for this indicator	9,410,269	20.6%
	Total area	45,698,163	100%



#### Natural landcover in floodplains

This indicator measures the amount of natural landcover in the estimated floodplain of rivers and streams within each catchment. It assesses the stream channel and its surrounding riparian buffer, measuring the percent of unaltered habitat like forests, wetlands, or open water (rather than agriculture or development). Intact vegetated buffers within the floodplain of rivers and streams provide aquatic habitat, improve water quality, reduce erosion and flooding, recharge groundwater, and more. This indicator originates from the National Land Cover Database and applies to the Environmental Protection Agency's estimated floodplain, which spatially defines areas estimated to be inundated by a 100-year flood (also known as the 1% annual chance flood).





#### Percent natural landcover within the estimated floodplain, by catchment

>90% natural landcover

>80-90% natural landcover

>70-80% natural landcover

>60-70% natural landcover

≤60% natural landcover

Not identified as a floodplain

Table 20: Indicator values for natural landcover in floodplains within Florida. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.

	Indicator Values: Percent natural landcover within the estimated floodplain, by catchment	Acres	Percent of Area	
↑ High	>90% natural landcover	11,951,129	26.2%	↑ In good
	>80-90% natural landcover	748,147	1.6%	condition
	>70-80% natural landcover	200,620	0.4%	↓ Not in good
	>60-70% natural landcover	154,221	0.3%	condition
	≤60% natural landcover	395,342	0.9%	
↓ Low	Not identified as a floodplain	22,869,978	50.0%	
	Area not evaluated for this indicator	9,378,726	20.5%	
	Total area	45,698,163	100%	



This indicator depicts the number of connected stream size classes in a river network between dams or waterfalls. River networks with a variety of connected stream classes help retain aquatic biodiversity in a changing climate by allowing species to access climate refugia and move between habitats. This indicator originates from the Southeast Aquatic Resources Partnership and applies to the Environmental Protection Agency's estimated floodplain, which spatially defines areas estimated to be inundated by a 100-year flood (also known as the 1% annual chance flood).





Table 21: Indicator values for network complexity within Florida. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.

	Indicator Values: Number of connected stream size classes	Acres	Percent of Area	
↑ High	7 size classes	24,730	<0.1%	
	6 size classes	2,511,351	5.5%	
	5 size classes	2,491,872	5.5%	
	4 size classes	2,791,383	6.1%	↑ In good condition
	3 size classes	1,758,480	3.8%	↓ Not in good
	2 size classes	2,072,503	4.5%	condition
	1 size class	1,695,050	3.7%	
↓ Low	Not identified as a floodplain	22,910,197	50.1%	
	Area not evaluated for this indicator	9,442,598	20.7%	
	Total area	45,698,163	100%	



This indicator measures the average percent of non-impervious cover within each catchment. High levels of impervious surface degrade water quality and alter freshwater flow, impacting both aquatic species communities and ecosystem services for people, like the availability of clean drinking water. This indicator originates from the National Land Cover Database.





#### Percent of catchment permeable

- >95% permeable (likely high water quality and supporting most sensitive aquatic species)
- >90-95% permeable (likely declining water quality and supporting most aquatic species)
- >70-90% permeable (likely degraded water quality and not supporting many aquatic species)
- ≤70% permeable (likely degraded instream flow, water quality, and aquatic species communities)

Table 22: Indicator values for permeable surface within Florida. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.

	Indicator Values: Percent of catchment permeable	Acres	Percent of Area	
↑ High	>95% permeable (likely high water quality and supporting most sensitive aquatic species)	28,484,809	62.3%	↑ In good condition
	>90-95% permeable (likely declining water quality and supporting most aquatic species)	2,401,914	5.3%	↓ Not in good condition
	>70-90% permeable (likely degraded water quality and not supporting many aquatic species)	3,781,797	8.3%	
↓ Low	≤70% permeable (likely degraded instream flow, water quality, and aquatic species communities)	1,640,128	3.6%	
	Area not evaluated for this indicator	9,389,514	20.5%	
	Total area	45,698,163	100%	

## Coastal & marine Atlantic coral & hardbottom

This indicator predicts the presence of coral and hardbottom in the Atlantic ocean based on direct observations, known locations of human-created structures like artificial reefs, and predictive models. The models use hardbottom observations and a suite of environmental predictors including measures of depth, seafloor topography and substrate, oceanography, and geography. Hardbottom provides an anchor for important seafloor habitats such as deep-sea corals, plants, and sponges, providing valuable structure that supports a wide range of invertebrate and fish species. This indicator combines data from The Nature Conservancy's South Atlantic Bight Marine Assessment and multiple National Oceanic and Atmospheric Administration datasets (deep-sea coral observations, shipwrecks, artificial reefs, and two projects predicting hardbottom distribution).




Table 23: Indicator values for Atlantic coral & hardbottom within Florida. A good condition threshold is not yet defined for this indicator.

	Indicator Values	Acres	Percent of Area
↑ High	Confirmed coral	323	<0.1%
	Confirmed natural or human-created hardbottom (shipwrecks, artificial reefs)	387,386	0.8%
	Predicted cold water coral mounds (Blake Plateau)	0	0%
	Highest probability of hardbottom (>80th percentile)	2,456	<0.1%
	High probability of hardbottom(>60th-80th percentile)	24,828	<0.1%
	Medium probability of hardbottom (>40th-60th percentile)	124,943	0.3%
	Low probability of hardbottom (>20th-40th percentile)	537,663	1.2%
	Lowest probability of hardbottom (≤20th percentile)	1,476,430	3.2%
↓ Low	Not identified as hardbottom	16,969,742	37.1%
	Area not evaluated for this indicator	26,174,393	57.3%
	Total area	45,698,163	100%

## Coastal & marine

#### Atlantic deep-sea coral richness

This indicator measures the number of deep-sea coral genera predicted to occur in the Atlantic Ocean at depths of approximately 50 m or below. It is based on coral observations and a suite of environmental predictors including measures of depth, seafloor topography and substrate, oceanography, and geography. This indicator combines probability models for 24 deep-sea coral genera to predict overall richness. Deep-sea corals provide valuable habitat structure that supports a wide range of invertebrate and fish species, and higher coral diversity typically creates more complex habitats occupied by more species. This indicator originates from a National Oceanic and Atmospheric Administration project characterizing the spatial distributions of deep-sea corals and hardbottom habitats in the U.S. Southeast Atlantic.




Table 24: Indicator values for Atlantic deep-sea coral richness within Florida. A good condition threshold is not yet defined for this indicator.

	Indicator Values	Acres	Percent of Area
↑ High	Highest predicted genus richness (>4)	0	0%
	High predicted genus richness (>3-4)	0	0%
	Medium predicted genus richness (>2-3)	0	0%
	Low predicted genus richness (>1-2)	0	0%
↓ Low	Lowest predicted genus richness (0-1)	598	<0.1%
	Area not evaluated for this indicator	45,697,565	100.0%
	Total area	45,698,163	100%

### Coastal & marine

#### Atlantic estuarine fish habitat

This indicator measures the condition of estuarine fish habitat along the Atlantic coast using metrics of water quality, marsh edges, seagrass and oyster reefs, fragmentation, human development, and more. Areas of excellent fish habitat are already in good condition and face few threats. Restoration opportunity areas are doing well in some respects, but restoration projects could significantly improve them. Degraded areas of opportunity face many challenges, and restoration projects are unlikely to increase available fish habitat unless particularly large in scope and scale. This indicator originates from the Atlantic Coast Fish Habitat Partnership's fish habitat conservation area mapping and prioritization project.






Table 25: Indicator values for Atlantic estuarine fish habitat within Florida. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.

	Indicator Values	Acres	Percent of Area	
↑ High	Final score of 80 (areas of excellent fish habitat)	0	0%	
	Final score of 70 (areas of excellent fish habitat)	6,175	<0.1%	↑ In good condition
	Final score of 60 (restoration opportunity areas)	137,625	0.3%	↓ Not in good condition
	Final score of 50 (restoration opportunity areas)	666,723	1.5%	
	Final score of 40 (restoration opportunity areas)	1,085,645	2.4%	
	Final score of 30 (restoration opportunity areas)	645,095	1.4%	
	Final score of 20 (restoration opportunity areas)	212,164	0.5%	
	Final score of 10 (degraded areas of opportunity)	41,987	<0.1%	
↓ Low	Final score of 0 (degraded areas of opportunity)	3,461	<0.1%	
	Area not evaluated for this indicator	42,899,288	93.9%	
	Total area	45,698,163	100%	



This indicator identifies important areas in the Atlantic Ocean for birds that feed exclusively or mainly at sea. It uses seasonal predictions of relative abundance for 19 species of marine birds (Audubon's shearwater, band-rumped storm petrel, black-capped petrel, black scoter, Bonaparte's gull, bridled tern, brown pelican, common loon, common tern, Cory's shearwater, great shearwater, Manx shearwater, Northern gannet, parasitic jaeger, red-throated loon, royal tern, sooty shearwater, sooty tern, white-winged scoter) based on sightings from boat-based surveys and marine environmental data like fronts, primary productivity, and ocean currents. This indicator originates from Duke University's Marine-life Data and Analysis Team marine bird models.





Land

>10th-20th percentile ≤10th percentile


Table 26: Indicator values for Atlantic marine birds within Florida. A good condition threshold is not yet defined for this indicator.

	Indicator Values: Percentile of importance for marine bird index species (across the full East Coast study area)	Acres	Percent of Area
↑ High	>90th percentile	902,103	2.0%
	>80th-90th percentile	209,827	0.5%
	>70th-80th percentile	52,682	0.1%
	>60th-70th percentile	32,096	<0.1%
	>50th-60th percentile	35,461	<0.1%
	>40th-50th percentile	14,001	<0.1%
	>30th-40th percentile	989	<0.1%
	>20th-30th percentile	0	0%
	>10th-20th percentile	0	0%
	≤10th percentile	0	0%
↓ Low	Land	12,825	<0.1%
	Area not evaluated for this indicator	44,438,178	97.2%
	Total area	45,698,163	100%



#### Atlantic marine mammals

This indicator identifies important areas in the Atlantic Ocean for dolphins, whales, and seals. It incorporates density predictions for 20 marine mammals species or species groups (Atlantic spotted dolphin, Atlantic white-sided dolphin, Clymene dolphin, common bottlenose dolphin, Cuvier's beaked whale, dwarf and pygmy sperm whales, fin whale, harbor porpoise, humpback whale, mesoplodont beaked whales, North Atlantic right whale, pantropical spotted dolphin, pilot whales, Risso's dolphin, rough-toothed dolphin, seals, short-beaked common dolphin, sperm whale, striped dolphin, unidentified beaked whales) based on sightings from boat-based and aerial surveys and data on oceanographic conditions. It uses marine mammal models developed by the Duke Marine Lab.





Percentile of importance for marine mammal index species (across the full East Coast study area)

>90th percentile

>80th-90th percentile

>70th-80th percentile

>70th-80th percentile
>60th-70th percentile
>50th-60th percentile
>40th-50th percentile
>30th-40th percentile
>20th-30th percentile
>10th-20th percentile
≤10th percentile
Land

Table 27: Indicator values for Atlantic marine mammals within Florida. A good condition threshold is not yet defined for this indicator.

	Indicator Values: Percentile of importance for marine mammal index species (across the full East Coast study area)	Acres	Percent of Area
↑ High	>90th percentile	510,074	1.1%
	>80th-90th percentile	234,188	0.5%
	>70th-80th percentile	61,705	0.1%
	>60th-70th percentile	13,109	<0.1%
	>50th-60th percentile	5,238	<0.1%
	>40th-50th percentile	14,013	<0.1%
	>30th-40th percentile	6,206	<0.1%
	>20th-30th percentile	50,775	0.1%
	>10th-20th percentile	57,683	0.1%
	≤10th percentile	47,392	0.1%
↓ Low	Land	149,780	0.3%
	Area not evaluated for this indicator	44,547,999	97.5%
	Total area	45,698,163	100%

## (Vy)

Coastal & marine

#### Coastal shoreline condition

This indicator assesses shoreline condition based on the presence of hardened structures like jetties, groins, and riprap, as well as other human development. By restricting the natural movement of sediment, shoreline armoring increases erosion, prevents the inland migration of coastal ecosystems in response to sea-level rise, and degrades habitat for birds, sea turtles, fish, plants, and other species both on and offshore. Natural shorelines in harder-to-develop coastal areas receive the highest shoreline condition scores, while hardened shorelines receive the lowest scores. This indicator originates from the National Oceanic and Atmospheric Administration's Environmental Sensitivity Index dataset.




Table 28: Indicator values for coastal shoreline condition within Florida. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.

	Indicator Values	Acres	Percent of Area	
↑ High	Natural and harder to develop	45,534	<0.1%	
	Natural	231,129	0.5%	↑ In good condition
	Partially armored and harder to develop	306	<0.1%	↓ Not in good condition
	Partially armored	5,768	<0.1%	
↓ Low	Armored	55,014	0.1%	
	Area not evaluated for this indicator	45,360,411	99.3%	
	Total area	45,698,163	100%	



#### **Estuarine coastal condition**

This indicator combines measures of water quality, sediment quality, contaminants in fish tissue, and benthic community condition to create an overall index of coastal estuarine condition. Estuaries serve as important nursery habitat for wildlife, including many species of fish and shellfish eaten as seafood. They also improve water quality by filtering out sediments and pollutants, provide recreational opportunities, and support coastal economies. This indicator originates from the Environmental Protection Agency's National Coastal Condition Assessment data.




Table 29: Indicator values for estuarine coastal condition within Florida. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.

	Indicator Values	Acres	Percent of Area	
↑ High	Good	818,226	1.8%	
	Fair to good	1,211,496	2.7%	↑ In good condition
	Fair	3,851,904	8.4%	↓ Not in good
↓ Low	Poor to fair	69,338	0.2%	condition
	Poor	7,377	<0.1%	
	Shallow estuary not assessed for condition	898,101	2.0%	
	Area not evaluated for this indicator	38,841,720	85.0%	
	Total area	45,698,163	100%	



# This indicator predicts the presence of coral and hardbottom in the Gulf of Mexico based on direct observations, acoustic surveys, designated Coral Habitat Areas of Particular Concern, and known locations of human-created structures like artificial reefs. Hardbottom provides an anchor for important seafloor habitats such as deep-sea corals, plants, and sponges, providing valuable structure that supports a wide range of invertebrate and fish species. Hardbottom is also sometimes associated with diverse chemosynthetic communities supported by bacteria that feed off of hydrocarbon seeps. This indicator combines data from multiple sources, including Bureau of Ocean Energy Management seismic water bottom anomalies, usSEABED sediments, several National Oceanic and




Table 30: Indicator values for Gulf coral & hardbottom within Florida. A good condition threshold is not yet defined for this indicator.

	Indicator Values	Acres	Percent of Area
↑ High	Confirmed hardbottom-associated species (corals, patch reef, chemosynthetic communities, or other organisms)	286	<0.1%
	Confirmed human-created hardbottom (shipwrecks, artificial reefs, decommissioned oil and gas platforms)	31,633	<0.1%
	Predicted hardbottom (fine resolution)	0	0%
	Coral Habitat Area of Particular Concern (HAPC)	1,400,140	3.1%
	Rock (coarse resolution)	136,485	0.3%
	Gravel (coarse resolution)	855,328	1.9%
↓ Low	Not identified as hardbottom	28,889,880	63.2%
	Area not evaluated for this indicator	14,384,411	31.5%
	Total area	45,698,163	100%

## (Vy)

Coastal & marine

#### Gulf deep-sea coral richness

This indicator measures the number of different deep-sea coral genera predicted to occur in the Gulf of Mexico at depths of 50 m or below. It is based on coral observations and a suite of environmental predictors including measures of depth, seafloor topography and substrate, oceanography, and geography. This indicator combines probability models for 28 deep-sea coral genera to predict overall richness. Deep-sea corals provide valuable habitat structure that supports a wide range of invertebrate and fish species, and higher coral diversity typically creates more complex habitats occupied by more species. This indicator originates from a National Oceanic and Atmospheric Administration project characterizing the spatial distributions of deep-sea corals and chemosynthetic communities in the U.S. Gulf of Mexico.




Table 31: Indicator values for Gulf deep-sea coral richness within Florida. A good condition threshold is not yet defined for this indicator.


	Indicator Values	Acres	Percent of Area
↑ High	Highest predicted average genus richness (>7)	3,151	<0.1%
	High predicted average genus richness (>5-7)	0	0%
	Medium predicted average genus richness (>3-5)	0	0%
↓ Low	Low predicted average genus richness (>1-3)	0	0%
	Lowest predicted average genus richness (0-1)	0	0%
	Area not evaluated for this indicator	45,695,012	100.0%
	Total area	45,698,163	100%



Coastal & marine

#### **Gulf marine mammals**

This indicator identifies important areas in the Gulf of Mexico for dolphins and whales. It incorporates monthly density predictions for 13 marine mammal species or species groups (Atlantic spotted dolphin, beaked whales, blackfish [which includes killer whale, melon-headed whale, false killer whale, pygmy killer whale], bottlenose dolphin, Bryde's whale, clymene dolphin, pantropical spotted dolphin, pilot whales, pygmy/dwarf sperm whales, Rice's whale, Risso's dolphin, sperm whale, spinner dolphin) based on sightings from boat-based and aerial surveys and data on oceanographic conditions. It uses marine mammal models developed by the National Oceanic and Atmospheric Administration as part of the Gulf of Mexico Marine Assessment Program for Protected Species (GoMMAPPS).





#### Percentile of importance for marine mammal index species (across larger analysis area)

>90th percentile

>80th-90th percentile

>70th-80th percentile

>60th-70th percentile

>50th-60th percentile

>40th-50th percentile

>30th-40th percentile

>20th-30th percentile

>10th-20th percentile

≤10th percentile


Land

Table 32: Indicator values for Gulf marine mammals within Florida. A good condition threshold is not yet defined for this indicator.

	Indicator Values: Percentile of importance for marine mammal index species (across larger analysis area)	Acres	Percent of Area
↑ High	>90th percentile	74,963	0.2%
	>80th-90th percentile	279,951	0.6%
	>70th-80th percentile	457,612	1.0%
	>60th-70th percentile	435,027	1.0%
	>50th-60th percentile	258,226	0.6%
	>40th-50th percentile	447,157	1.0%
	>30th-40th percentile	302,118	0.7%
	>20th-30th percentile	565,389	1.2%
	>10th-20th percentile	409,003	0.9%
	≤10th percentile	4,508,002	9.9%
↓ Low	Land	135,680	0.3%
	Area not evaluated for this indicator	37,825,033	82.8%
	Total area	45,698,163	100%



This indicator identifies important areas in the Gulf of Mexico for sea turtles. It incorporates monthly density predictions for four species (green, Kemp's ridley, leatherback, and loggerhead sea turtles) based on sightings from boat-based and aerial surveys and data on oceanographic conditions. It uses sea turtle models developed by the National Oceanic and Atmospheric Administration as part of the Gulf of Mexico Marine Assessment Program for Protected Species (GoMMAPPS).






Table 33: Indicator values for Gulf sea turtles within Florida. A good condition threshold is not yet defined for this indicator.

	Indicator Values: Percentile of importance for sea turtle index species (across larger analysis area)	Acres	Percent of Area
↑ High	>90th percentile	4,947,181	10.8%
	>80th-90th percentile	1,334,687	2.9%
	>70th-80th percentile	171,595	0.4%
	>60th-70th percentile	8,797	<0.1%
↓ Low	≤60th percentile	1,555,412	3.4%
	Land	2,305,459	5.0%
	Area not evaluated for this indicator	35,375,032	77.4%
	Total area	45,698,163	100%



This indicator represents important habitat for coastal island-dependent species across the Southeast. Because the isolation of islands can make them ecologically unique and protect them from disturbance and mainland predators, they often serve as important habitat for many species of mammals, plants, and insects, as well as breeding coastal birds and sea turtles. The highest scores go to island critical habitat for six threatened and endangered animal and plant species: piping plover, loggerhead sea turtle, Cape Sable thoroughwort, Florida semaphore cactus, silver rice rat, and Bartram's hairstreak butterfly. This indicator uses U.S. Fish and Wildlife Service critical habitat data and island boundaries from the U.S. Geological Survey and Esri.





Table 34: Indicator values for island habitat within Florida. A good condition threshold is not yet defined for this indicator.

	Indicator Values	Acres	Percent of Area
↑ High	Island critical habitat for any of six threatened and endangered species (piping plover, loggerhead sea turtle, Cape Sable thoroughwort, Florida semaphore cactus, silver rice rat, or Bartram's hairstreak butterfly)	23,647	<0.1%
	Other island area	413,836	0.9%
↓ Low	Not a coastal island	29,998,712	65.6%
	Area not evaluated for this indicator	15,261,968	33.4%
	Total area	45,698,163	100%

## Coastal & marine Marine hic

#### Marine highly migratory fish

This indicator identifies important foraging and spawning areas for highly migratory fish in the Atlantic Ocean and Gulf of Mexico. It uses physical capture and satellite tag observations, remote sensing of environmental variables, and physical oceanographic data to analyze the habitat preferences of three species (skipjack tuna, bluefin tuna, and blue shark) at various life stages. It originates from European Commission Joint Research Centre global fish models.





#### Percentile of importance for bluefin and skipjack tuna or blue shark

- >90th percentile
- >80th-90th percentile
- >70th-80th percentile
- >60th-70th percentile
  - >50th-60th percentile
  - >40th-50th percentile
  - >30th-40th percentile
- ≤30th percentile

Table 35: Indicator values for marine highly migratory fish within Florida. A good condition threshold is not yet defined for this indicator.

	Indicator Values: Percentile of importance for bluefin and skipjack tuna or blue shark	Acres	Percent of Area
↑ High	>90th percentile	158,302	0.3%
	>80th-90th percentile	115,950	0.3%
	>70th-80th percentile	108,496	0.2%
	>60th-70th percentile	146,752	0.3%
	>50th-60th percentile	108,488	0.2%
	>40th-50th percentile	270,822	0.6%
	>30th-40th percentile	2,328,641	5.1%
↓ Low	≤30th percentile	110,962	0.2%
	Area not evaluated for this indicator	42,349,750	92.7%
	Total area	45,698,163	100%

# Coastal & marine Resilient coastal sites

This indicator depicts the capacity of coastal habitats to migrate to adjacent lowlands in order to sustain biodiversity and natural services under increasing inundation from sea-level rise. It is based on the physical and condition characteristics of current tidal complexes, their predicted migration space, and surrounding buffer areas. These characteristics include marsh complex size, shared edge with migration space, sediment balance, water quality, natural landcover, landform diversity, and more. This indicator originates from The Nature Conservancy's Resilient Coastal Sites project.




Table 36: Indicator values for resilient coastal sites within Florida. A good condition threshold is not yet defined for this indicator.

	Indicator Values	Acres	Percent of Area
↑ High	Most resilient	0	0%
	More resilient	2,768,618	6.1%
	Slightly more resilient	258,699	0.6%
	Average/median resilience	349,437	0.8%
	Slightly less resilient	19,317	<0.1%
↓ Low	Less resilient	7,934	<0.1%
	Least resilient	21,751	<0.1%
	Area not evaluated for this indicator	42,272,405	92.5%
	Total area	45,698,163	100%



This indicator represents the presence of seagrass in the Atlantic Ocean and Gulf of Mexico. Seagrasses provide food and habitat for a range of marine and estuarine wildlife, including fish, sea turtles, shrimp, crabs, oysters, and more. They also produce oxygen, filter water, control erosion, and buffer storms. Seagrasses serve as an important indicator of the overall health of coastal ecosystems because they are sensitive to water quality and require sufficiently clear water for sunlight to penetrate. This indicator originates from the National Oceanic and Atmospheric Administration's Marine Cadastre.





Table 37: Indicator values for seagrass within Florida. A good condition threshold is not yet defined for this indicator.

	Indicator Values	Acres	Percent of Area
↑ High	Seagrass present	2,289,118	5.0%
	Area not evaluated for this indicator	43,409,045	95.0%
	Total area	45,698,163	100%



## South Atlantic beach birds

This indicator is an index of habitat suitability for four shorebird species (American oystercatcher, Wilson's plover, least tern, piping plover) in the South Atlantic, based on observed abundance. It assesses beaches and nearby onshore habitats. Shorebirds' relative use of beaches and neighboring habitats for nesting, foraging, and breeding is an indicator of ecosystem health and quality. This indicator combines bird data from the U.S. Geological Survey and state waterbird biologists in FL, GA, SC, and NC.





#### Percentile of importance for beach bird index species

- >80th percentile
  - >60th-80th percentile
- >40th-60th percentile
- >20th-40th percentile
  - ≤20th percentile
- 320th percentite
  - Open water or not identified as a priority

Table 38: Indicator values for South Atlantic beach birds within Florida. A good condition threshold is not yet defined for this indicator.

	Indicator Values: Percentile of importance for beach bird index species	Acres	Percent of Area
↑ High	>80th percentile	969	<0.1%
	>60th-80th percentile	701	<0.1%
	>40th-60th percentile	0	0%
↓ Low	>20th-40th percentile	0	0%
	≤20th percentile	161	<0.1%
	Open water or not identified as a priority	246,784	0.5%
	Area not evaluated for this indicator	45,449,548	99.5%
	Total area	45,698,163	100%



## South Atlantic maritime forest

This indicator depicts the maritime forest currently present in the South Atlantic. Since maritime forest has been substantially reduced from its historic extent, protecting the remaining acreage is particularly important. This ecosystem supports a unique suite of plants that tolerate wind, salt, and flooding, as well as many species of birds, mammals, and reptiles. It also helps buffer the coastline from storms. This indicator originates from LANDFIRE landcover.




Table 39: Indicator values for South Atlantic maritime forest within Florida. A good condition threshold is not yet defined for this indicator.

	Indicator Values	Acres	Percent of Area
↑ High	Maritime forest	10,148	<0.1%
↓ Low	Not identified as maritime forest	3,076,392	6.7%
	Area not evaluated for this indicator	42,611,623	93.2%
	Total area	45,698,163	100%



This indicator uses remote sensing to calculate the unvegetated-vegetated ratio of tidal wetlands, which compares how much of a wetland is not covered by plants (e.g., sediment, rocks, open water) to how much is covered by plants. Marshes that maintain a higher proportion of vegetation tend to be more stable and resilient to threats like sea-level rise, erosion, and coastal development. This ratio, and how it changes over time, is a good surrogate for salt marsh degradation processes like sediment loss and conversion to open water. This indicator originates from a U.S. Geological Survey project on an unvegetated to vegetated ratio for coastal wetlands.




Table 40: Indicator values for stable coastal wetlands within Florida. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.

	Indicator Values	Acres	Percent of Area	
↑ High	Stable coastal wetlands	1,108,435	2.4%	
	Other coastal wetlands	279,743	0.6%	↑ In good condition
↓ Low	Not identified as coastal wetlands	10,622,311	23.2%	↓ Not in good condition
	Area not evaluated for this indicator	33,687,673	73.7%	
	Total area	45,698,163	100%	

## **Threats**

### Sea-level rise

NOAA's sea-level rise (SLR) inundation models represent areas likely to experience flooding at high tide based on each foot of SLR above current levels. Darker blue areas will experience flooding first, and at greater depth, compared to lighter blue areas. These models are not linked to a future timeframe; see the projections below. NOAA calculates the inundation footprint at "mean higher high water", or the average highest daily tide. The area covered in each SLR scenario includes areas projected to be inundated at lower levels. For example, the area inundated by 4 ft of SLR also includes areas inundated by 3 ft, 2 ft, 1 ft, and 0 ft of SLR (where 0 ft represents current levels).

To explore additional SLR information, please see NOAA's <u>Sea Level Rise Viewer</u>.




Table 41: Extent of flooding by projected average highest daily tide due to sea level rise within Florida. Values from the NOAA sea-level rise inundation data.

Feet of sea-level rise	Acres	Percent of Area
0 feet	6,792,251	14.9%
1 foot	7,575,392	16.6%
2 feet	8,053,294	17.6%
3 feet	8,422,260	18.4%
4 feet	8,764,845	19.2%
5 feet	9,222,145	20.2%
6 feet	9,708,184	21.2%
7 feet	10,216,613	22.4%
8 feet	10,847,358	23.7%
9 feet	11,464,988	25.1%
10 feet	11,913,006	26.1%
Not projected to be inundated by up to 10 feet	16,659,224	36.5%
Sea-level rise unlikely to be a threat (inland counties)	13,551,898	29.7%
Sea-level rise data unavailable	3,574,034	7.8%
Total area	45,698,163	100%


Table 42: Projected sea level rise by decade within Florida. Values are based on area-weighted averages of decadal projections for 1-degree grid cells that overlap this area based on <u>NOAA's 2022 Sea Level Rise</u>

<u>Report.</u> 2060 corresponds to the <u>SECAS goal</u>: a 10% or greater improvement in the health, function, and connectivity of Southeastern ecosystems by 2060.

SLR Scenario	2020 (ft)	2030 (ft)	2040 (ft)	2050 (ft)	2060 (ft)	2070 (ft)	2080 (ft)	2090 (ft)	2100 (ft)
Low	0.31	0.48	0.66	0.82	0.97	1.1	1.2	1.3	1.4
Intermediate- low	0.34	0.54	0.75	0.96	1.2	1.4	1.6	1.8	2
Intermediate	0.35	0.56	0.81	1.1	1.4	1.8	2.3	2.9	3.6
Intermediate- high	0.35	0.59	0.9	1.3	1.8	2.5	3.3	4.2	5.1
High	0.35	0.62	0.98	1.5	2.2	3.2	4.3	5.5	6.8

## **Urban growth**

The FUTURES urban growth model predicts the likelihood that an area will urbanize at every decade from 2020 to 2100. Developed areas from the 2021 National Landcover Database serve as the baseline for current urban areas. The model simulates landscape change based on trends in population growth, local development suitability factors, and an urban patch-growing algorithm. It considers environmental drivers like distance to floodplain, slope, and available infrastructure, and even socio-economic status. The probability of urbanization for each area reflects how many times it urbanized out of 50 model runs.



12.6% of this area is already urban in 2021, and an additional 12.3% has at least a moderate probability of urbanizing by 2060.

Table 43: Extent of projected urbanization by decade within Florida. Values from <u>FUTURES model</u> <u>projections for the contiguous United States</u> developed by the <u>Center for Geospatial Analytics</u>, NC State University. 2060 corresponds to the <u>SECAS goal</u>: a 10% or greater improvement in the health, function, and connectivity of Southeastern ecosystems by 2060.

Decade	Acres	Percent of Area
Urban in 2021	5,764,045	12.6%
2030 projected extent	5,965,613	13.1%
2040 projected extent	6,107,901	13.4%
2050 projected extent	6,219,537	13.6%
2060 projected extent	6,315,061	13.8%
2070 projected extent	6,396,501	14.0%
2080 projected extent	6,453,162	14.1%
2090 projected extent	6,486,677	14.2%
2100 projected extent	6,502,683	14.2%
Not projected to urbanize by 2100	29,938,055	65.5%
Total area	45,698,163	100%

# **Ownership and Partners**

# Conserved lands ownership





Table 44: Extent of ownership class within Florida. Protected areas are derived from the <u>Protected Areas</u> <u>Database of the United States</u> (PAD-US v3.0) and include Fee, Designation, Easement, Marine, and Proclamation (Dept. of Defense lands only) boundaries. Note: areas are based on the polygon boundary of this area compared to protected area polygons, rather than pixel-level analyses used elsewhere in this report. Also note: PAD-US v3.0 includes protected areas that may overlap within a given area; this may cause the area within and between the following categories to be greater than the actual ground area.

Ownership	Acres	Percent of Area
Federal	4,053,137	8.9%
State/province	4,081,260	8.9%
Regional	1,701,923	3.7%
Local	553,203	1.2%
Joint	1,201	<0.1%
Private non-profit conserved lands	74,185	0.2%
Private conservation land	1,087,114	2.4%
Tribal	46,199	0.1%
Designation	10,496,950	23.0%
Ownership unknown	36,251	<0.1%

## **Land protection status**





- Managed for biodiversity (disturbance events proceed or are mimicked)
- Managed for biodiversity (disturbance events suppressed)
- Managed for multiple uses (subject to extractive uses such as mining or logging, or OHV use)
- No known mandate for biodiversity protection

Table 45: Extent of land protection status within Florida. Protected areas are derived from the <u>Protected Areas Database of the United States</u> (PAD-US v3.0) and include Fee, Designation, Easement, Marine, and Proclamation (Dept. of Defense lands only) boundaries. Note: areas are based on the polygon boundary of this area compared to protected area polygons, rather than pixel-level analyses used elsewhere in this report. Also note: PAD-US v3.0 includes protected areas that may overlap within a given area; this may cause the area within and between the following categories to be greater than the actual ground area.

Land Protection Status	Acres	Percent of Area
Managed for biodiversity (disturbance events proceed or are mimicked)	3,885,994	8.5%
Managed for biodiversity (disturbance events suppressed)	9,833,424	21.5%
Managed for multiple uses (subject to extractive uses such as mining or logging, or OHV use)	6,370,492	13.9%
No known mandate for biodiversity protection	2,041,516	4.5%

### **Protected Areas**

- Everglades National Park (Unknown; 1,538,630 acres)
- EVER (NPS; 1,532,380 acres)
- Florida Keys National Marine Sanctuary (Unknown; 1,403,774 acres)
- Marjory Stoneman Douglas Wilderness (1,338,506 acres)
- National Forests in Florida (USDA FOREST SERVICE; 1,203,406 acres)
- BICY (NPS; 683,630 acres)
- Big Bend Seagrasses Aquatic Preserve (Unknown; 680,739 acres)
- Everglades and Francis S. Taylor Wildlife Management Area (Trustees of the Internal Improvement Trust Fund; 669,448 acres)
- Eglin Air Force Base (452,797 acres)
- Lake Okeechobee (440,465 acres)
- Florida Keys Areas to be Avoided (Unknown; 390,961 acres)
- Pinellas County Aquatic Preserve (Unknown; 351,938 acres)
- Apalachicola National Estuarine Research Reserve (Unknown; 235,675 acres)
- Blackwater River State Forest (Trustees of the Internal Improvement Trust Fund; 206,191 acres)
- Tate's Hell State Forest (Trustees of the Internal Improvement Trust Fund; 204,575 acres)
- Great White Heron National Wildlife Refuge (Trustees of the Internal Improvement Trust Fund; 203,810 acres)

- Key West National Wildlife Refuge (Trustees of the Internal Improvement Trust Fund; 188,974 acres)
- Key West National Wildlife Refuge (Unknown; 187,836 acres)
- Withlacoochee State Forest (Trustees of the Internal Improvement Trust Fund; 165,562 acres)
- Biscayne National Park (Unknown; 157,164 acres)
- BISC (NPS; 155,762 acres)
- Great White Heron National Wildlife Refuge (Unknown; 134,231 acres)
- Merritt Island National Wildlife Refuge (Unknown; 129,419 acres)
- Biscayne Bay-Card Sound Spiny Lobster Sanctuary (Unknown; 125,475 acres)
- Apalachicola National Estuarine Research Reserve (Trustees of the Internal Improvement Trust Fund; 114,615 acres)
- ... and 8,486 more protected areas ...

## **Nearby land trusts**

Click here to search for land trusts within 500 miles of this area on the Land Trust Alliance website.

## **Credits**

This report was generated by the Southeast Conservation Blueprint Explorer, which was developed by <u>Astute Spruce, LLC</u> in partnership with the U.S. Fish and Wildlife Service under the <u>Southeast Conservation Adaptation Strategy</u>.

#### **Data credits**

Land ownership and conservation status is derived from the <u>Protected Areas Database of the United States</u> (PAD-US v3.0).

Future urban growth estimates derived from <u>FUTURES model projections for the contiguous United States</u> developed by the <u>Center for Geospatial Analytics</u>, NC State University.

Sea level rise data are derived from the National Oceanic and Atmospheric Administration's <u>Sea Level Rise Inundation Depth Data</u> and the <u>2022 Sea Level Rise Technical Report</u>.