The Southeast Blueprint Indicators

The Southeast Conservation Blueprint is based on 61 natural and cultural resource indicators that represent terrestrial, freshwater, and coastal/marine ecosystems.



Continental

15 Southeast states + U.S. Atlantic and Gulf waters



Terrestrial

- **East Coastal Plain open pine birds:** Identifies areas within the longleaf pine range east of the Mississippi River where open pine habitat management would most benefit 6 bird species (*source: East Gulf Coastal Plain Joint Venture JV*).
- Equitable access to potential parks: Prioritizes places to create new parks to fill gaps in access to green space within socially vulnerable communities (source: Trust for Public Land ParkServe, Centers for Disease Control Social Vulnerability Index).
- **Fire frequency:** Estimates the number of times an area has been burned from 2013-2021 (*source: Landsat Burned Areas, Monitoring Trends in Burn Severity*).
- **Great Plains perennial grasslands:** Evaluates grassland condition using the percent of perennial forbs and grasses (*source: Rangeland Analysis Program RAP*).
- **Greenways & trails:** Assesses the natural condition and connected length of recreational paths (*source: OpenStreetMap, National Land Cover Database NLCD*).
- **Intact habitat cores:** Measures the size of large, unfragmented patches of natural habitat (*source: Esri* green infrastructure data).
- Interior Southeast grasslands: Index of known grasslands, areas with potentially compatible management, and grassland geology (*source: RAP*).
- Mississippi Alluvial Valley forest birds protection: Prioritizes new areas for protection to benefit 3 forest breeding bird species that need large interior cores of bottomland hardwood habitat (source: Lower Mississippi Valley Joint Venture - LMVJV).
- Mississippi Alluvial Valley forest birds reforestation: Prioritizes areas for reforestation to benefit 3 forest breeding bird species that need large interior cores of bottomland hardwood habitat (source: LMVJV).

- **Playas:** Depicts the location and condition of playas in the western Great Plains (*source: Playa Lakes JV*).
- **Resilient terrestrial sites:** Index depicting the ability of terrestrial sites to continue supporting biodiversity and ecosystem function in the face of climate change (*source: The Nature Conservancy TNC Resilient Land*).
- South Atlantic amphibian & reptile areas: Identifies important herpetofauna habitat designated as Priority Amphibian and Reptile Conservation Areas (PARCAs) (source: South Atlantic PARCAs).
- **South Atlantic forest birds:** Evaluates habitat suitability for 12 upland hardwood and forested wetland bird species (*source: Southeast Gap Analysis Program GAP, Designing Sustainable Landscapes*).
- South Atlantic low-urban historic landscapes: Assesses the degree of urban development surrounding sites on the National Register of Historic Places (source: National Park Service - NPS, State Historic Preservation Offices - SHPOs).
- **Urban park size:** Measures the size of parks larger than 5 acres in the urban environment (*source: Protected Areas Database of the United States - PAD-US, NLCD, Census urban areas).*
- West Coastal Plain & Ouachitas forested wetland birds: Evaluates habitat suitability within bottomland hardwood forests and riparian areas for 5 bird species (*source: LMVJV*).
- West Coastal Plain & Ouachitas open pine birds: Identifies areas with existing pine trees that, if managed for open condition, could support a population of 3 pine bird species (*source: LMVJV*).
- West Gulf Coast mottled duck nesting: Depicts coastal marshes and grasslands in TX and LA that are important for mottled duck nesting (source: Gulf Coast Prairie Landscape Conservation Cooperative).



Continental

15 Southeast states + U.S. Atlantic and Gulf waters



- Atlantic migratory fish habitat: Assesses migratory fish habitat condition based on water quality, connectivity, flow alteration, and more (*source: Atlantic Coast Fish Habitat Partnership ACFHP*).
- **Gulf migratory fish connectivity:** Depicts how far upstream migratory fish species have been observed (source: TNC Southeast Aquatic Connectivity Assessment Project, U.S. Fish & Wildlife Service - USFWS critical habitat).
- Imperiled aquatic species: Counts the number of aquatic Species of Greatest Conservation Need observed in each watershed (source: Southeast Aquatic Resources Partnership HUC12 species summaries, Environmental Protection Agency - EPA estimated floodplain).
- Natural landcover in floodplains: Measures the percent natural landcover in the estimated floodplain by catchment (source: NLCD, National Hydrography Dataset Plus high resolution NHDPlus HR, EPA estimated floodplain).

- **Network complexity:** Counts the number of connected stream size classes in a river network between dams or waterfalls (*source: SARP network complexity data, NHDPlus HR, EPA estimated floodplain).*
- **Permeable surface:** Measures the average percent of non-impervious landcover within each catchment (*source: NLCD, NHDPlus HR*).
- West Virginia imperiled aquatic species: Counts the number of globally imperiled, threatened, or endangered aquatic animal species observed in each watershed (source: EPA EnviroAtlas HUC12 species summaries, EPA estimated floodplain).



Continental

15 Southeast states + U.S. Atlantic and Gulf waters



- Atlantic coral & hardbottom: Predicts the presence of coral and hardbottom habitat based on direct observations, known human-created structures, and distribution models (source: TNC South Atlantic Bight Marine Assessment; National Oceanic & Atmospheric Administration - NOAA deep-sea coral observations, shipwrecks, artificial reefs, hardbottom distribution data).
- Atlantic deep-sea coral richness: Measures the number of deep-sea coral genera predicted to occur offshore (source: NOAA deep-sea coral distribution data).
- Atlantic estuarine fish habitat: Assesses estuarine fish habitat condition based on water quality, marsh edges, seagrass, oyster reefs, and more (*source: ACFHP*).
- Atlantic marine birds: Identifies important areas for 19 bird species that feed exclusively or mainly at sea (source: Duke Marine-life Data & Analysis Team).
- Atlantic marine mammals: Identifies important areas for 20 species and species groups of dolphins, whales, and seals (source: Duke Marine Lab).
- **Coastal shoreline condition:** Evaluates shoreline condition based on the presence of hardened structures and ease of development in coastal areas (source: NOAA Environmental Sensitivity Index ESI, Coastal Barrier Resource System).
- **Estuarine coastal condition:** Assesses estuarine water quality, sediment quality, contaminants in fish tissue, and benthic community condition (*source: EPA National Coastal Condition Assessment*).
- **Gulf coral & hardbottom:** Predicts the presence of coral and hardbottom habitat based on direct observations, acoustic surveys, designated coral habitat, and known human-created structures (source: Bureau of Ocean Energy Management seismic water bottom anomalies; usSEABED sediments; coral Habitat Areas of Particular Concern; Dept. of Homeland Security oil and gas platforms; NOAA deep-sea coral observations, shipwrecks, artificial reefs).

- **Gulf deep-sea coral richness:** Measures the number of deep-sea coral genera predicted to occur offshore (*source: NOAA deep-sea coral distribution data*).
- **Gulf marine mammals:** Identifies important areas for 13 species or species groups of dolphins and whales (source: NOAA Gulf of Mexico Marine Assessment Program for Protected Species - GoMMAPPS).
- **Gulf sea turtles:** Identifies important areas for 4 species of sea turtles (*source: NOAA GoMMAPPS*).
- **Island habitat:** Identifies important habitat for islanddependent species (*source: USFWS critical habitat, Esri/U.S. Geological Survey - USGS island boundaries*).
- **Marine highly migratory fish:** Identifies important foraging and spawning areas for 3 highly migratory fish species at various life stages (*source: European Commission global fish models*).
- **Resilient coastal sites:** Assesses the ability of coastal sites to migrate inland to sustain biodiversity and natural services under increasing inundation from sealevel rise (*source: TNC Resilient Coastal Sites*).
- **Seagrass:** Identifies seagrass locations (*source: NOAA Marine Cadastre*).
- **South Atlantic beach birds:** Predicts habitat suitability for 4 shorebird species (*source: bird data from USGS and state waterbird biologists*).
- **South Atlantic maritime forest:** Identifies maritime forest locations (*source: LANDFIRE EVT*).
- **Stable coastal wetlands:** Evaluates marsh stability and resilience based on the ratio of plants to non-plants (e.g., open water or sediment) *(source: USGS unvegetated-vegetated ratio data).*

Caribbean



Puerto Rico & the U.S. Virgin Islands + nearshore marine waters



Terrestrial

- **Caribbean greenways & trails:** Assesses the natural condition and connected length of recreational paths (source: OpenStreetMap, NOAA Coastal Change Analysis Program C-CAP).
- **Caribbean habitat patch size (large islands):** Measures the size of natural, unfragmented habitat patches on large islands (*source: LANDFIRE Existing Vegetation Type - EVT and roads*).
- **Caribbean habitat patch size (small islands):** Measures the size of natural, unfragmented habitat patches on small islands (*source: LANDFIRE EVT and roads*).
- **Caribbean island habitat:** Assesses the importance of island habitat for federally listed and other imperiled species based on the presence of imperiled and invasive animals (*source: USFWS critical habitat, Puerto Rico GAP, U.S. Virgin Islands GAP, Island Conservation Threatened Island Biodiversity Database).*
- **Caribbean karst habitat:** Identifies natural karst areas with limited human alteration (*source: LANDFIRE EVT; karst layers from USGS, NPS, Puerto Rico Dept. of Natural & Environmental Resources*).
- **Caribbean landscape condition:** Assesses the degree of human alteration of natural habitat at multiple scales, from a specific site to the surrounding landscape (*source: LANDFIRE EVT*).
- **Caribbean low-urban historic landscapes:** Assesses the degree of urban development surrounding historic sites (source: LANDFIRE EVT; historic sites from OpenStreetMap, Puerto Rico SHPO, University of the Virgin Islands).
- **Caribbean reforestation potential:** Prioritizes areas to increase tree cover based on current land uses and benefits to drinking water supplies (*source: LANDFIRE EVT*).
- **Caribbean urban park size:** Measures the size of parks in the urban environment (*source: Census urban areas; protected areas from PAD-US, OpenStreetMap, Puerto Rico Protected Natural Areas, TNC*).



- **Caribbean natural landcover in floodplains:** Measures the percent of natural landcover in the estimated floodplain by catchment (*source: LANDFIRE EVT, NHDPlus HR, Federal Emergency Management Agency - FEMA National Flood Hazard Layer*).
- **Caribbean network complexity:** Counts the number of connected stream size classes in a river network between dams or waterfalls (*source: SARP network complexity data, NHDPlus HR, FEMA National Flood Hazard Layer*).
- **Caribbean permeable surface:** Measures the average percent of non-impervious landcover in each catchment (*source: NOAA C-CAP, NHDPlus HR*).



Coastal & Marine

- **Caribbean beach habitat:** Evaluates beach habitat for 6 species of beach-nesting birds and sea turtles (source: PR & USVI GAP, State of the World's Sea Turtles; beaches from OpenStreetMap, NOAA Caribbean Climate Adaptation Program).
- **Caribbean coastal shoreline condition:** Assesses the extent of coastal shoreline armoring (*source: NOAA Continuously Updated Shoreline Product*).
- **Caribbean fish hotspots:** Predicts levels of fish density and diversity based on the proximity of mangroves, seagrass, and coral (*source: NOAA C-CAP, TNC benthic habitat map*).
- **Caribbean fish nursery habitat:** Identifies fish nursery, spawning, and concentration areas like lagoons, bays, estuaries, and reefs (*source: NOAA ESI*).
- **Caribbean seagrass:** Identifies the presence and density of seagrass at various depths (*source: TNC benthic habitat map, NOAA bathymetry*).
- **Caribbean shallow hardbottom & coral:** Identifies the presence of hardbottom habitat and coral and assesses coral resilience to climate change (*source: TNC benthic habitat map and coral climate refugia*).