Southeast Conservation Blueprint Summary

for Virginia

Created 01/19/2024

Table of Contents

About the Southeast Blueprint	3
Southeast Blueprint Priorities	4
Hubs and Corridors	6
Indicator Summary	8
Threats	64
Ownership and Partners	68
Credits	73

The Southeast Conservation Adaptation Strategy



Yer The Southeast Conservation Blueprint 2023

[THIS PAGE INTENTIONALLY LEFT BLANK]

About the Southeast Blueprint

The Southeast Conservation Blueprint is the primary product of the <u>Southeast Conservation Adaptation</u> <u>Strategy</u> (SECAS). It is a living, spatial plan to achieve the SECAS vision of a connected network of lands and waters across the Southeast and Caribbean. The Blueprint is regularly updated to incorporate new data, partner input, and information about on-the-ground conditions.

The Blueprint identifies priority areas based on a suite of natural and cultural resource indicators representing terrestrial, freshwater, and marine ecosystems. A connectivity analysis identifies corridors that link coastal and inland areas and span climate gradients.

For more information:

- Visit the <u>Blueprint webpage</u>
- Review the <u>Blueprint 2023 Development Process</u>
- View and download the Blueprint data and make maps on the Blueprint page of the SECAS Atlas

We're here to help!

- Do you have a question about the Blueprint?
- Would you like help using the Blueprint to support a proposal or inform a decision?
- Do you have a suggestion on how to improve the Blueprint? The Blueprint and its inputs are regularly revised based on input from people like you.
- Do you have feedback on how to improve the Simple Viewer interface?

If you need help or have questions, <u>contact Southeast Blueprint staff</u> by reaching out to a member of the user support team.

We're here to support you. We really mean it. It's what we do!

Southeast Blueprint Priorities



Priorities for a connected network of lands and waters

- Highest priority
- High priority
- Medium priority
- Priority connections

Priority Categories

For a connected network of lands and waters

In total, Blueprint priorities and priority connections cover roughly 50% of the Southeast Blueprint geography.

Highest priority

Areas where conservation action would make the biggest impact, based on a suite of natural and cultural resource indicators. This class covers roughly 10% of the Southeast Blueprint geography.

High priority

Areas where conservation action would make a big impact, based on a suite of natural and cultural resource indicators. This class covers roughly 15% of the Southeast Blueprint geography.

Medium priority

Areas where conservation action would make an above-average impact, based on a suite of natural and cultural resource indicators. This class covers roughly 20% of the Southeast Blueprint geography.

Priority connections

Connections between priority areas that cover the shortest distance possible while routing through as much Blueprint priority as possible. This class covers roughly 5% of the Southeast Blueprint geography.

Priority Category	Acres	Percent of Area
Highest priority	2,231,311	8.2%
High priority	3,586,962	13.1%
Medium priority	5,892,933	21.5%
Priority connections	1,282,457	4.7%
Lower priority	14,366,617	52.5%
Total area	27,360,280	100%

Table 1: Extent of each Blueprint priority category within Virginia.

Hubs and Corridors

The Blueprint uses a least-cost path connectivity analysis to identify corridors that link hubs across the shortest distance possible, while also routing through as much Blueprint priority as possible.

Inland hubs are large patches (~5,000+ acres) of highest priority Blueprint areas and/or protected lands, connected by inland corridors. Marine and estuarine hubs are large estuaries and large patches (~5,000+ acres) of highest priority Blueprint areas. Marine and estuarine corridors connect those hubs within broad marine mammal movement areas.





- Inland continental hubs
- Inland continental corridors
- Marine & estuarine continental hubs
- Marine & estuarine continental corridors
- Not a hub or corridor

Table 2: Extent of hubs and corridors within Virginia.

Туре	Acres	Percent of Area
Inland continental hubs	3,382,929	12.4%
Inland continental corridors	3,514,200	12.8%
Marine & estuarine continental hubs	1,378,254	5.0%
Marine & estuarine continental corridors	228,165	0.8%
Not a hub or corridor	18,856,734	68.9%
Total area	27,360,280	100%

Indicator Summary

Table 3: Terrestrial indicators.

Indicator	Present
East Coastal Plain open pine birds	\checkmark
Equitable access to potential parks	\checkmark
Fire frequency	\checkmark
<u>Greenways & trails</u>	\checkmark
Intact habitat cores	\checkmark
Interior Southeast grasslands	\checkmark
Resilient terrestrial sites	\checkmark
South Atlantic amphibian & reptile areas	\checkmark
South Atlantic forest birds	\checkmark
South Atlantic low-urban historic landscapes	\checkmark
<u>Urban park size</u>	\checkmark

Table 4: Freshwater indicators.

Indicator	Present
Atlantic migratory fish habitat	\checkmark
Imperiled aquatic species	\checkmark
Natural landcover in floodplains	\checkmark
Network complexity	\checkmark
Permeable surface	\checkmark
West Virginia imperiled aquatic species	-

Table 5: Coastal & marine indicators.

Indicator	Present
Atlantic coral & hardbottom	\checkmark
Atlantic deep-sea coral richness	-
Atlantic estuarine fish habitat	\checkmark
Atlantic marine birds	\checkmark
Atlantic marine mammals	\checkmark
Coastal shoreline condition	\checkmark
Estuarine coastal condition	\checkmark
Gulf coral & hardbottom	-
Island habitat	\checkmark
Marine highly migratory fish	-
Resilient coastal sites	\checkmark
Seagrass	\checkmark
South Atlantic beach birds	-
South Atlantic maritime forest	\checkmark
Stable coastal wetlands	\checkmark



This indicator identifies areas within the historic longleaf pine range east of the Mississippi River where creating or maintaining open pine habitat would most benefit six focal species of birds (Bachman's sparrow, red-cockaded woodpecker, Henslow's sparrow, red-headed woodpecker, Northern bobwhite, brown-headed nuthatch). It prioritizes areas for open pine conservation based on suitability for longleaf pine, feasibility of prescribed burning, proximity to protected lands, habitat suitability for focal bird species, and proximity to bird source populations. It originates from the East Gulf Coastal Plain Joint Venture's prioritization of areas for open pine ecosystem restoration.





Priority for open pine conservation for focal bird species

- High priority (score >80-100)
- Medium-high priority (score >60-80)
- Medium priority (score >40-60)
- Medium-low priority (score >20-40)
- Low priority (score 0-20)
- Not a priority (not identified as upland pine)

Table 6: Indicator values for east coastal plain open pine birds within Virginia. A good condition threshold is not yet defined for this indicator.

	Indicator Values: Priority for open pine conservation for focal bird species	Acres	Percent of Area
↑ High	High priority (score >80-100)	0	0%
	Medium-high priority (score >60-80)	0	0%
	Medium priority (score >40-60)	0	0%
	Medium-low priority (score >20-40)	2,485	<0.1%
↓ Low	Low priority (score 0-20)	331,608	1.2%
	Not a priority (not identified as upland pine)	793,216	2.9%
	Area not evaluated for this indicator	26,232,971	95.9%
	Total area	27,360,280	100%



This cultural resource indicator prioritizes places to create new parks that would fill gaps in equitable access to open space within socially vulnerable communities in urban areas. It identifies areas where residents currently lack access to parks within a 10-minute walk (accounting for walkable road networks and access barriers like highways and fences), then prioritizes based on park need using demographic and environmental metrics. Parks help improve public health, foster a conservation ethic by providing opportunities for people to connect with nature, and support critical ecosystem services. This indicator originates from the Trust for Public Land's ParkServe park priority areas and the Center for Disease Control's Social Vulnerability Index.





Priority for a new park that would create nearby equitable access

- Very high priority
- High priority
- Moderate priority
 - Not identified as a priority (within urban areas)

Table 7: Indicator values for equitable access to potential parks within Virginia. A good condition threshold is not yet defined for this indicator.

	Indicator Values: Priority for a new park that would create nearby equitable access	Acres	Percent of Area
↑ High	Very high priority	67,144	0.2%
	High priority	69,395	0.3%
↓ Low	Moderate priority	57,990	0.2%
	Not identified as a priority (within urban areas)	25,706,086	94.0%
	Area not evaluated for this indicator	1,459,666	5.3%
	Total area	27,360,280	100%



This indicator uses remote sensing to estimate the number of times an area has been burned from 2013 to 2021. Many Southeastern ecosystems rely on regular, low-intensity fires to maintain habitat, encourage native plant growth, and reduce wildfire risk. This indicator combines burned area layers from U.S. Geological Survey Landsat data and the inter-agency Monitoring Trends in Burn Severity program. Landsat-based fire predictions within the range of longleaf pine are also available through <u>Southeast FireMap</u>.





Burned 3+ times from 2013-2021
Burned 2 times from 2013-2021
Burned 1 time from 2013-2021
Not burned from 2013-2021 or row crop

Table 8: Indicator values for fire frequency within Virginia. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.

	Indicator Values	Acres	Percent of Area	
↑ High	Burned 3+ times from 2013-2021	6,546	<0.1%	
	Burned 2 times from 2013-2021	22,485	<0.1%	↑ In good condition
	Burned 1 time from 2013-2021	219,875	0.8%	↓ Not in good
↓ Low	Not burned from 2013-2021 or row crop	27,099,148	99.0%	condition
	Area not evaluated for this indicator	12,227	<0.1%	
	Total area	27,360,280	100%	



This cultural resource indicator measures both the natural condition and connected length of greenways and trails to characterize the quality of the recreational experience. Natural condition is based on the amount of impervious surface surrounding the path. Connected length captures how far a person can go without leaving a dedicated path, based on common distances for walking, running, and biking. This indicator originates from OpenStreetMap data and the National Land Cover Database.





- Mostly natural and connected for ≥40 km
- Mostly natural and connected for 5 to <40 km or partly natural and connected for ≥40 km
- Mostly natural and connected for 1.9 to <5 km, partly natural and connected for 5 to <40 km, or developed and connected for ≥40 km
- Mostly natural and connected for <1.9 km, partly natural and connected for 1.9 to <5 km, or developed and connected for 5 to <40 km
- Partly natural and connected for <1.9 km or developed and connected for 1.9 to <5 km
- Developed and connected for <1.9 km</p>
- Sidewalk
- Not identified as a trail, sidewalk, or other path

Table 9: Indicator values for greenways & trails within Virginia. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.

	Indicator Values	Acres	Percent of Area	
↑ High	Mostly natural and connected for ≥40 km	23,805	<0.1%	
	Mostly natural and connected for 5 to <40 km or partly natural and connected for ≥40 km	27,280	<0.1%	
	Mostly natural and connected for 1.9 to <5 km, partly natural and connected for 5 to <40 km, or developed and connected for ≥40 km	30,428	0.1%	
	Mostly natural and connected for <1.9 km, partly natural and connected for 1.9 to <5 km, or developed and connected for 5 to <40 km	14,111	<0.1%	↑ In good condition
	Partly natural and connected for <1.9 km or developed and connected for 1.9 to <5 km	7,415	<0.1%	↓ Not in good condition
	Developed and connected for <1.9 km	11,489	<0.1%	
	Sidewalk	45,357	0.2%	
↓ Low	Not identified as a trail, sidewalk, or other path	27,192,162	99.4%	
	Area not evaluated for this indicator	8,232	<0.1%	
	Total area	27,360,280	100%	



This indicator represents the size of large, unfragmented patches of natural habitat. It identifies minimally disturbed natural areas at least 100 acres in size and greater than 200 meters wide. Large areas of intact natural habitat are important for many wildlife species, including reptiles and amphibians, birds, and large mammals. This indicator originates from Esri's green infrastructure data.



Basemap credits: © <u>Mapbox</u> © <u>OpenStreetMap Improve this basemap</u>



Large core (>10,000 acres) Medium core (>1,000-10,000 acres) Small core (>100-1,000 acres) Not a core

108

54

Table 10: Indicator values for intact habitat cores within Virginia. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.

	Indicator Values	Acres	Percent of Area	
↑ High	Large core (>10,000 acres)	2,852,686	10.4%	
	Medium core (>1,000-10,000 acres)	6,596,377	24.1%	
	Small core (>100-1,000 acres)	3,078,133	11.3%	↑ In good condition
↓ Low	Not a core	14,824,853	54.2%	↓ Not in good condition
	Area not evaluated for this indicator	8,232	<0.1%	
	Total area	27,360,280	100%	



This indicator represents grasslands in the interior southeastern United States, which support important plants, birds, and pollinators. It includes grasslands with and without trees that are historically maintained by geology (e.g., outcrops, glades, and barrens), fire (e.g., Piedmont prairies), and/or the regular violent flooding on the banks of high-energy rivers known as "riverscour" (e.g., riverscour prairies). Known grasslands receive the highest scores, followed by bumble bee habitat buffers around known sites, areas in potentially compatible management, and restoration opportunities within grassland geology. This indicator combines data from multiple sources, including the Southeastern Grasslands Institute, Central Hardwoods Joint Venture, Rangeland Analysis Platform, and more.





Known grassland

Known grassland buffer

Potentially compatible management within grassland geology (undeveloped powerline right-of-way or perennial forbs and grasses) Potentially compatible management outside of grassland geology (undeveloped powerline right-of-way or perennial forbs and grasses) Grassland geology

Grassland less likely

Table 11: Indicator values for Interior Southeast grasslands within Virginia. A good condition threshold is not yet defined for this indicator.

	Indicator Values	Acres	Percent of Area
↑ High	Known grassland	6,701	<0.1%
	Known grassland buffer	44,399	0.2%
	Potentially compatible management within grassland geology (undeveloped powerline right-of-way or perennial forbs and grasses)	1,103,099	4.0%
	Potentially compatible management outside of grassland geology (undeveloped powerline right-of-way or perennial forbs and grasses)	2,578,260	9.4%
	Grassland geology	4,212,982	15.4%
↓ Low	Grassland less likely	12,086,525	44.2%
	Area not evaluated for this indicator	7,328,315	26.8%
	Total area	27,360,280	100%



This indicator depicts an area's capacity to maintain species diversity and ecosystem function in the face of climate change. It measures two factors that influence resilience. The first, landscape diversity, reflects the number of microhabitats and climatic gradients created by topography, elevation, and hydrology. The second, local connectedness, reflects the degree of habitat fragmentation and strength of barriers to species movement. Highly resilient sites contain many different habitat niches that support biodiversity, and allow species to move freely through the landscape to find suitable microclimates as the climate changes. This indicator originates from The Nature Conservancy's Resilient Land data.





Most resilient
More resilient
Slightly more resilient
Average/median resilience
Slightly less resilient
Less resilient
Least resilient
Developed

Table 12: Indicator values for resilient terrestrial sites within Virginia. A good condition threshold is not yet defined for this indicator.

	Indicator Values	Acres	Percent of Area
↑ High	Most resilient	425,146	1.6%
	More resilient	4,140,228	15.1%
	Slightly more resilient	4,813,424	17.6%
	Average/median resilience	5,952,687	21.8%
	Slightly less resilient	2,788,092	10.2%
	Less resilient	2,711,616	9.9%
	Least resilient	634,428	2.3%
↓ Low	Developed	3,303,975	12.1%
	Area not evaluated for this indicator	2,590,684	9.5%
	Total area	27,360,280	100%



This indicator represents Priority Amphibian and Reptile Conservation Areas (PARCAs) in the South Atlantic. PARCA is an expert-driven, nonregulatory designation that includes places capable of supporting viable amphibian and reptile populations, places occupied by rare or imperiled species, and places rich in biodiversity or species unique to that geographic area (i.e., endemism). Reptiles and amphibians are a critical part of the Southeast region's rich biodiversity and many populations are declining in the face of threats like habitat loss, invasive species, and climate change.





Priority Amphibian and Reptile Conservation Area (PARCA) Not a Priority Amphibian and Reptile Conservation Area (PARCA) Table 13: Indicator values for South Atlantic amphibian & reptile areas within Virginia. A good condition threshold is not yet defined for this indicator.

	Indicator Values	Acres	Percent of Area
↑ High	Priority Amphibian and Reptile Conservation Area (PARCA)	226,436	0.8%
↓ Low	Not a Priority Amphibian and Reptile Conservation Area (PARCA)	6,956,605	25.4%
	Area not evaluated for this indicator	20,177,240	73.7%
	Total area	27,360,280	100%

Priority Amphibian and Reptile Conservation Areas:

Fairystone

No description available for PARCAs in Virginia

Fort Pickett

No description available for PARCAs in Virginia

Great Dismal Swamp

No description available for PARCAs in Virginia

North Landing

No description available for PARCAs in Virginia

South Mountains

The rugged landscape of the South Mountains chain, combined with cool and clear streams, create a unique topographical oasis in the western Piedmont of North Carolina known as the South Mountains PARCA. Elevations rising up to 3,000 feet provide habitat for both the timber rattlesnake and the narrowly endemic South Mountain gray-cheeked salamander.



This indicator is an index of habitat suitability for twelve upland hardwood and forested wetland bird species (wood thrush, whip-poor-will, American woodcock, red-headed woodpecker, Chuck-will's widow, hooded warbler, Kentucky warbler, Acadian flycatcher, Northern parula, black-throated green warbler, prothonotary warbler, Swainson's warbler) based on patch size and other ecosystem characteristics such as proximity to water and proximity to forest and ecotone edge. The needs of these species are increasingly restrictive at higher index values, reflecting better quality habitat. It originates from Southeast Gap Analysis Program and Designing Sustainable Landscapes bird habitat models.





- Very large patches near water (potential for presence of Swainson's warbler)
- Large patches often near water (potential for presence of Northern parula, black-throated green warbler, or prothonotary warbler)
 Medium patches (potential for presence of Acadian flycatcher)
 Small patches often near water (potential for presence of hooded warbler or Kentucky warbler)
- Very small patches or near open areas (potential for presence of wood thrush, whip-poor-will, red-headed woodpecker, Chuck-will's widow, or American woodcock)
- Less potential for presence of forest bird index species

Table 14: Indicator values for South Atlantic forest birds within Virginia. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.

	Indicator Values	Acres	Percent of Area	
↑ High	Very large patches near water (potential for presence of Swainson's warbler)	269,148	1.0%	
	Large patches often near water (potential for presence of Northern parula, black-throated green warbler, or prothonotary warbler)	623,824	2.3%	
	Medium patches (potential for presence of Acadian flycatcher)	1,850,219	6.8%	
	Small patches often near water (potential for presence of hooded warbler or Kentucky warbler)	352,952	1.3%	↑ In good condition
	Very small patches or near open areas (potential for presence of wood thrush, whip- poor-will, red-headed woodpecker, Chuck- will's widow, or American woodcock)	3,201,834	11.7%	↓ Not in good condition
↓ Low	Less potential for presence of forest bird index species	877,465	3.2%	
	Area not evaluated for this indicator	20,184,839	73.8%	
	Total area	27,360,280	100%	

South Atlantic low-urban historic landscapes

This cultural resource indicator is an index of sites on the National Register of Historic Places surrounded by limited urban development. It identifies significant historic places that remain connected to their context in the natural world. It uses the National Land Cover Database and historic places data from the National Park Service and various state historic resource agencies.





Historic place with nearby low-urban buffer Historic place with nearby high-urban buffer Not in the National Register of Historic Places Table 15: Indicator values for South Atlantic low-urban historic landscapes within Virginia. A good condition threshold is not yet defined for this indicator.

	Indicator Values	Acres	Percent of Area
↑ High	Historic place with nearby low-urban buffer	29,176	0.1%
	Historic place with nearby high-urban buffer	5,112	<0.1%
↓ Low	Not in the National Register of Historic Places	6,335,966	23.2%
	Area not evaluated for this indicator	20,990,026	76.7%
	Total area	27,360,280	100%



This cultural resource indicator measures the size of parks larger than 5 acres in the urban environment. Protected natural areas in urban environments provide urban residents a nearby place to connect with nature, and offer refugia for some species. This indicator complements the equitable access to potential parks indicator by capturing the value of existing parks. It originates from the Protected Areas Database of the United States, Census urban areas, and the National Land Cover Database.





75+ acre urban park
50 to <75 acre urban park
30 to <50 acre urban park
10 to <30 acre urban park

- 5 to <10 acre urban park
- <5 acre urban park or not identified as an urban park

Table 16: Indicator values for urban park size within Virginia. A good condition threshold is not yet defined for this indicator.

	Indicator Values	Acres	Percent of Area
↑ High	75+ acre urban park	182,355	0.7%
	50 to <75 acre urban park	8,946	<0.1%
	30 to <50 acre urban park	8,910	<0.1%
↓ Low	10 to <30 acre urban park	16,550	<0.1%
	5 to <10 acre urban park	5,614	<0.1%
	<5 acre urban park or not identified as an urban park	27,137,465	99.2%
	Area not evaluated for this indicator	440	<0.1%
	Total area	27,360,280	100%



This indicator measures the condition of migratory fish habitat along the Atlantic coast within each catchment, using metrics of water quality, aquatic connectivity, habitat fragmentation, flow alteration, and more. Areas of excellent fish habitat are already in good condition and face few threats. Restoration opportunity areas are doing well in some respects, but restoration projects could significantly improve them. Degraded areas of opportunity face many challenges, and restoration projects are unlikely to increase available fish habitat unless particularly large in scope and scale. This indicator originates from the Atlantic Coast Fish Habitat Partnership's fish habitat conservation area mapping and prioritization project.





Final score of 80 (areas of excellent fish habitat) Final score of 70 (areas of excellent fish habitat) Final score of 60 (restoration opportunity areas) Final score of 50 (restoration opportunity areas) Final score of 40 (restoration opportunity areas) Final score of 30 (restoration opportunity areas) Final score of 20 (restoration opportunity areas) Final score of 10 (degraded areas of opportunity) Final score of 0 (degraded areas of opportunity) Table 17: Indicator values for Atlantic migratory fish habitat within Virginia. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.

	Indicator Values	Acres	Percent of Area	
↑ High	Final score of 80 (areas of excellent fish habitat)	161	<0.1%	
	Final score of 70 (areas of excellent fish habitat)	45,112	0.2%	↑ In good condition
	Final score of 60 (restoration opportunity areas)	402,213	1.5%	↓ Not in good condition
	Final score of 50 (restoration opportunity areas)	395,394	1.4%	
	Final score of 40 (restoration opportunity areas)	519,112	1.9%	
	Final score of 30 (restoration opportunity areas)	432,586	1.6%	
	Final score of 20 (restoration opportunity areas)	125,030	0.5%	
	Final score of 10 (degraded areas of opportunity)	21,036	<0.1%	
↓ Low	Final score of 0 (degraded areas of opportunity)	965	<0.1%	
	Area not evaluated for this indicator	25,418,671	92.9%	
	Total area	27,360,280	100%	



This indicator measures the number of aquatic animal Species of Greatest Conservation Need (SGCN) observed within each 12-digit HUC subwatershed, including fish, mussels, snails, crayfish, and amphibians. SGCN are identified in State Wildlife Action Plans as most in need of conservation action. This indicator captures patterns of rare and endemic aquatic species diversity. It originates from state Natural Heritage Program data collected by the Southeast Aquatic Resources Partnership and applies to the Environmental Protection Agency's estimated floodplain, which spatially defines areas estimated to be inundated by a 100-year flood (also known as the 1% annual chance flood).





Table 18: Indicator values for imperiled aquatic species within Virginia. A good condition threshold is not yet defined for this indicator.

	Indicator Values: Number of aquatic animal Species of Greatest Conservation Need observed	Acres	Percent of Area
↑ High	8+ species	36,521	0.1%
	7 species	7,437	<0.1%
	6 species	12,493	<0.1%
	5 species	26,307	<0.1%
	4 species	28,199	0.1%
	3 species	93,867	0.3%
	2 species	184,649	0.7%
	1 species	565,702	2.1%
	0 species	1,401,198	5.1%
↓ Low	Not identified as a floodplain (excluding West Virginia)	23,551,136	86.1%
	Area not evaluated for this indicator	1,452,770	5.3%
	Total area	27,360,280	100%



This indicator measures the amount of natural landcover in the estimated floodplain of rivers and streams within each catchment. It assesses the stream channel and its surrounding riparian buffer, measuring the percent of unaltered habitat like forests, wetlands, or open water (rather than agriculture or development). Intact vegetated buffers within the floodplain of rivers and streams provide aquatic habitat, improve water quality, reduce erosion and flooding, recharge groundwater, and more. This indicator originates from the National Land Cover Database and applies to the Environmental Protection Agency's estimated floodplain, which spatially defines areas estimated to be inundated by a 100-year flood (also known as the 1% annual chance flood).





Percent natural landcover within the estimated floodplain, by catchment

- >90% natural landcover >80-90% natural landcover
- >70-80% natural landcover
- >60-70% natural landcover
- ≤60% natural landcover
- Not identified as a floodplain
Table 19: Indicator values for natural landcover in floodplains within Virginia. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.

	Indicator Values: Percent natural landcover within the estimated floodplain, by catchment	Acres	Percent of Area	
↑ High	>90% natural landcover	1,460,897	5.3%	∧ In good
	>80-90% natural landcover	206,192	0.8%	condition
	>70-80% natural landcover	129,623	0.5%	↓ Not in good
	>60-70% natural landcover	107,871	0.4%	condition
	≤60% natural landcover	451,790	1.7%	
↓ Low	Not identified as a floodplain	23,551,462	86.1%	
	Area not evaluated for this indicator	1,452,444	5.3%	
	Total area	27,360,280	100%	



This indicator depicts the number of connected stream size classes in a river network between dams or waterfalls. River networks with a variety of connected stream classes help retain aquatic biodiversity in a changing climate by allowing species to access climate refugia and move between habitats. This indicator originates from the Southeast Aquatic Resources Partnership and applies to the Environmental Protection Agency's estimated floodplain, which spatially defines areas estimated to be inundated by a 100-year flood (also known as the 1% annual chance flood).





Number of connected stream size classes 7 size classes 6 size classes 5 size classes 4 size classes 3 size classes 2 size classes 1 size class Not identified as a floodplain

Table 20: Indicator values for network complexity within Virginia. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.

	Indicator Values: Number of connected stream size classes	Acres	Percent of Area	
↑ High	7 size classes	316,444	1.2%	
	6 size classes	413,684	1.5%	
	5 size classes	429,867	1.6%	
	4 size classes	482,245	1.8%	\uparrow In good condition
	3 size classes	285,835	1.0%	↓ Not in good
	2 size classes	297,659	1.1%	condition
	1 size class	123,412	0.5%	
\downarrow Low	Not identified as a floodplain	23,551,745	86.1%	
	Area not evaluated for this indicator	1,459,389	5.3%	
	Total area	27,360,280	100%	



This indicator measures the average percent of non-impervious cover within each catchment. High levels of impervious surface degrade water quality and alter freshwater flow, impacting both aquatic species communities and ecosystem services for people, like the availability of clean drinking water. This indicator originates from the National Land Cover Database.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap



Percent of catchment permeable

- >95% permeable (likely high water quality and supporting most sensitive aquatic species)
- >90-95% permeable (likely declining water quality and supporting most aquatic species)
- >70-90% permeable (likely degraded water quality and not supporting many aquatic species)

54

108

≤70% permeable (likely degraded instream flow, water quality, and aquatic species communities)

Table 21: Indicator values for permeable surface within Virginia. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.

	Indicator Values: Percent of catchment permeable	Acres	Percent of Area	
↑ High	>95% permeable (likely high water quality and supporting most sensitive aquatic species)	23,084,346	84.4%	↑ In good condition
	>90-95% permeable (likely declining water quality and supporting most aquatic species)	1,032,484	3.8%	↓ Not in good condition
	>70-90% permeable (likely degraded water quality and not supporting many aquatic species)	1,235,208	4.5%	
↓ Low	≤70% permeable (likely degraded instream flow, water quality, and aquatic species communities)	555,471	2.0%	
	Area not evaluated for this indicator	1,452,770	5.3%	
	Total area	27,360,280	100%	

Coastal & marine Atlantic coral & hardbottom

This indicator predicts the presence of coral and hardbottom in the Atlantic ocean based on direct observations, known locations of human-created structures like artificial reefs, and predictive models. The models use hardbottom observations and a suite of environmental predictors including measures of depth, seafloor topography and substrate, oceanography, and geography. Hardbottom provides an anchor for important seafloor habitats such as deep-sea corals, plants, and sponges, providing valuable structure that supports a wide range of invertebrate and fish species. This indicator combines data from The Nature Conservancy's South Atlantic Bight Marine Assessment and multiple National Oceanic and Atmospheric Administration datasets (deep-sea coral observations, shipwrecks, artificial reefs, and two projects predicting hardbottom distribution).







Confirmed coral

Confirmed natural or human-created hardbottom (shipwrecks, artificial reefs)

54

108

Predicted cold water coral mounds (Blake Plateau) Highest probability of hardbottom (>80th percentile) High probability of hardbottom(>60th-80th percentile) Medium probability of hardbottom (>40th-60th percentile) Low probability of hardbottom (>20th-40th percentile) Lowest probability of hardbottom (≤20th percentile) Not identified as hardbottom Table 22: Indicator values for Atlantic coral & hardbottom within Virginia. A good condition threshold is not yet defined for this indicator.

	Indicator Values	Acres	Percent of Area
↑ High	Confirmed coral	297	<0.1%
	Confirmed natural or human-created hardbottom (shipwrecks, artificial reefs)	6,279	<0.1%
	Predicted cold water coral mounds (Blake Plateau)	0	0%
	Highest probability of hardbottom (>80th percentile)	0	0%
	High probability of hardbottom(>60th-80th percentile)	0	0%
	Medium probability of hardbottom (>40th-60th percentile)	0	0%
	Low probability of hardbottom (>20th-40th percentile)	0	0%
	Lowest probability of hardbottom (≤20th percentile)	0	0%
↓ Low	Not identified as hardbottom	4,270,820	15.6%
	Area not evaluated for this indicator	23,082,885	84.4%
	Total area	27,360,280	100%



This indicator measures the condition of estuarine fish habitat along the Atlantic coast using metrics of water quality, marsh edges, seagrass and oyster reefs, fragmentation, human development, and more. Areas of excellent fish habitat are already in good condition and face few threats. Restoration opportunity areas are doing well in some respects, but restoration projects could significantly improve them. Degraded areas of opportunity face many challenges, and restoration projects are unlikely to increase available fish habitat unless particularly large in scope and scale. This indicator originates from the Atlantic Coast Fish Habitat Partnership's fish habitat conservation area mapping and prioritization project.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap



Final score of 80 (areas of excellent fish habitat) Final score of 70 (areas of excellent fish habitat) Final score of 60 (restoration opportunity areas) Final score of 50 (restoration opportunity areas) Final score of 40 (restoration opportunity areas) Final score of 30 (restoration opportunity areas) Final score of 20 (restoration opportunity areas) Final score of 10 (degraded areas of opportunity) Final score of 0 (degraded areas of opportunity)

108

54

Table 23: Indicator values for Atlantic estuarine fish habitat within Virginia. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.

	Indicator Values	Acres	Percent of Area	
↑ High	Final score of 80 (areas of excellent fish habitat)	0	0%	
	Final score of 70 (areas of excellent fish habitat)	13,095	<0.1%	↑ In good condition
	Final score of 60 (restoration opportunity areas)	108,958	0.4%	↓ Not in good condition
	Final score of 50 (restoration opportunity areas)	199,330	0.7%	
	Final score of 40 (restoration opportunity areas)	535,491	2.0%	
	Final score of 30 (restoration opportunity areas)	914,308	3.3%	
	Final score of 20 (restoration opportunity areas)	781,518	2.9%	
	Final score of 10 (degraded areas of opportunity)	175,580	0.6%	
↓ Low	Final score of 0 (degraded areas of opportunity)	42,011	0.2%	
	Area not evaluated for this indicator	24,589,988	89.9%	
	Total area	27,360,280	100%	



This indicator identifies important areas in the Atlantic Ocean for birds that feed exclusively or mainly at sea. It uses seasonal predictions of relative abundance for 19 species of marine birds (Audubon's shearwater, band-rumped storm petrel, black-capped petrel, black scoter, Bonaparte's gull, bridled tern, brown pelican, common loon, common tern, Cory's shearwater, great shearwater, Manx shearwater, Northern gannet, parasitic jaeger, red-throated loon, royal tern, sooty shearwater, sooty tern, white-winged scoter) based on sightings from boat-based surveys and marine environmental data like fronts, primary productivity, and ocean currents. This indicator originates from Duke University's Marine-life Data and Analysis Team marine bird models.





Percentile of importance for marine bird index species (across the full East Coast study area)

>90th percentile
>80th-90th percentile
>70th-80th percentile
>60th-70th percentile
>50th-60th percentile
>40th-50th percentile
>30th-40th percentile
>20th-30th percentile
>10th-20th percentile
≤10th percentile
Land

Table 24: Indicator values for Atlantic marine birds within Virginia. A good condition threshold is not yet defined for this indicator.

	Indicator Values: Percentile of importance for marine bird index species (across the full East Coast study area)	Acres	Percent of Area
↑ High	>90th percentile	215,050	0.8%
	>80th-90th percentile	16,543	<0.1%
	>70th-80th percentile	0	0%
	>60th-70th percentile	0	0%
	>50th-60th percentile	0	0%
	>40th-50th percentile	0	0%
	>30th-40th percentile	0	0%
	>20th-30th percentile	0	0%
	>10th-20th percentile	0	0%
	≤10th percentile	0	0%
↓ Low	Land	1,079	<0.1%
	Area not evaluated for this indicator	27,127,608	99.1%
	Total area	27,360,280	100%



This indicator identifies important areas in the Atlantic Ocean for dolphins, whales, and seals. It incorporates density predictions for 20 marine mammals species or species groups (Atlantic spotted dolphin, Atlantic whitesided dolphin, Clymene dolphin, common bottlenose dolphin, Cuvier's beaked whale, dwarf and pygmy sperm whales, fin whale, harbor porpoise, humpback whale, mesoplodont beaked whales, North Atlantic right whale, pantropical spotted dolphin, pilot whales, Risso's dolphin, rough-toothed dolphin, seals, short-beaked common dolphin, sperm whale, striped dolphin, unidentified beaked whales) based on sightings from boat-based and aerial surveys and data on oceanographic conditions. It uses marine mammal models developed by the Duke Marine Lab.









Percentile of importance for marine mammal index species (across the full East Coast study area)

>90th percentile >80th-90th percentile >70th-80th percentile >60th-70th percentile >50th-60th percentile >40th-50th percentile >30th-40th percentile >20th-30th percentile >10th-20th percentile ≤10th percentile Land

Table 25: Indicator values for Atlantic marine mammals within Virginia. A good condition threshold is not yet defined for this indicator.

	Indicator Values: Percentile of importance for marine mammal index species (across the full East Coast study area)	Acres	Percent of Area
↑ High	>90th percentile	205,848	0.8%
	>80th-90th percentile	64,976	0.2%
	>70th-80th percentile	33,948	0.1%
	>60th-70th percentile	6,152	<0.1%
	>50th-60th percentile	0	0%
	>40th-50th percentile	0	0%
	>30th-40th percentile	0	0%
	>20th-30th percentile	0	0%
	>10th-20th percentile	0	0%
	≤10th percentile	0	0%
↓ Low	Land	30,210	0.1%
	Area not evaluated for this indicator	27,019,146	98.8%
	Total area	27,360,280	100%



This indicator assesses shoreline condition based on the presence of hardened structures like jetties, groins, and riprap, as well as other human development. By restricting the natural movement of sediment, shoreline armoring increases erosion, prevents the inland migration of coastal ecosystems in response to sea-level rise, and degrades habitat for birds, sea turtles, fish, plants, and other species both on and offshore. Natural shorelines in harder-to-develop coastal areas receive the highest shoreline condition scores, while hardened shorelines receive the lowest scores. This indicator originates from the National Oceanic and Atmospheric Administration's Environmental Sensitivity Index dataset.





Natural and harder to develop

- Natural
- Partially armored and harder to develop
- Partially armored
- Armored

Table 26: Indicator values for coastal shoreline condition within Virginia. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.

	Indicator Values	Acres	Percent of Area	
↑ High	Natural and harder to develop	25,273	<0.1%	
	Natural	104,234	0.4%	↑ In good condition
	Partially armored and harder to develop	131	<0.1%	↓ Not in good condition
	Partially armored	2,555	<0.1%	
\downarrow Low	Armored	14,854	<0.1%	
	Area not evaluated for this indicator	27,213,233	99.5%	
	Total area	27,360,280	100%	

Coastal & marine Estuarine coastal condition

This indicator combines measures of water quality, sediment quality, contaminants in fish tissue, and benthic community condition to create an overall index of coastal estuarine condition. Estuaries serve as important nursery habitat for wildlife, including many species of fish and shellfish eaten as seafood. They also improve water quality by filtering out sediments and pollutants, provide recreational opportunities, and support coastal economies. This indicator originates from the Environmental Protection Agency's National Coastal Condition Assessment data.





Table 27: Indicator values for estuarine coastal condition within Virginia. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.

	Indicator Values	Acres	Percent of Area	
↑ High	Good	371,002	1.4%	
	Fair to good	155,082	0.6%	↑ In good condition
	Fair	307,193	1.1%	↓ Not in good
↓ Low	Poor to fair	3,010	<0.1%	condition
	Poor	107,389	0.4%	
	Shallow estuary not assessed for condition	396,221	1.4%	
	Area not evaluated for this indicator	26,020,383	95.1%	
	Total area	27,360,280	100%	

Coastal & marine

This indicator represents important habitat for coastal island-dependent species across the Southeast. Because the isolation of islands can make them ecologically unique and protect them from disturbance and mainland predators, they often serve as important habitat for many species of mammals, plants, and insects, as well as breeding coastal birds and sea turtles. The highest scores go to island critical habitat for six threatened and endangered animal and plant species: piping plover, loggerhead sea turtle, Cape Sable thoroughwort, Florida semaphore cactus, silver rice rat, and Bartram's hairstreak butterfly. This indicator uses U.S. Fish and Wildlife Service critical habitat data and island boundaries from the U.S. Geological Survey and Esri.





Island critical habitat for any of six threatened and endangered species (piping plover, loggerhead sea turtle, Cape Sable thoroughwort, Florida semaphore cactus, silver rice rat, or Bartram's hairstreak butterfly)

- Other island area
- Not a coastal island

Table 28: Indicator values for island habitat within Virginia. A good condition threshold is not yet defined for this indicator.

	Indicator Values	Acres	Percent of Area
↑ High	Island critical habitat for any of six threatened and endangered species (piping plover, loggerhead sea turtle, Cape Sable thoroughwort, Florida semaphore cactus, silver rice rat, or Bartram's hairstreak butterfly)	0	0%
	Other island area	417,382	1.5%
↓ Low	Not a coastal island	7,179,679	26.2%
	Area not evaluated for this indicator	19,763,219	72.2%
	Total area	27,360,280	100%



This indicator depicts the capacity of coastal habitats to migrate to adjacent lowlands in order to sustain biodiversity and natural services under increasing inundation from sea-level rise. It is based on the physical and condition characteristics of current tidal complexes, their predicted migration space, and surrounding buffer areas. These characteristics include marsh complex size, shared edge with migration space, sediment balance, water quality, natural landcover, landform diversity, and more. This indicator originates from The Nature Conservancy's Resilient Coastal Sites project.





Table 29: Indicator values for resilient coastal sites within Virginia. A good condition threshold is not yet defined for this indicator.

	Indicator Values	Acres	Percent of Area
↑ High	Most resilient	69,385	0.3%
	More resilient	46,540	0.2%
	Slightly more resilient	30,965	0.1%
	Average/median resilience	50,552	0.2%
	Slightly less resilient	3,477	<0.1%
	Less resilient	2,039	<0.1%
↓ Low	Least resilient	12,307	<0.1%
	Area not evaluated for this indicator	27,145,016	99.2%
	Total area	27,360,280	100%



This indicator represents the presence of seagrass in the Atlantic Ocean and Gulf of Mexico. Seagrasses provide food and habitat for a range of marine and estuarine wildlife, including fish, sea turtles, shrimp, crabs, oysters, and more. They also produce oxygen, filter water, control erosion, and buffer storms. Seagrasses serve as an important indicator of the overall health of coastal ecosystems because they are sensitive to water quality and require sufficiently clear water for sunlight to penetrate. This indicator originates from the National Oceanic and Atmospheric Administration's Marine Cadastre.





Seagrass present

Table 30: Indicator values for seagrass within Virginia. A good condition threshold is not yet defined for this indicator.

	Indicator Values	Acres	Percent of Area
↑ High	Seagrass present	41,268	0.2%
	Area not evaluated for this indicator	27,319,012	99.8%
	Total area	27,360,280	100%



This indicator depicts the maritime forest currently present in the South Atlantic. Since maritime forest has been substantially reduced from its historic extent, protecting the remaining acreage is particularly important. This ecosystem supports a unique suite of plants that tolerate wind, salt, and flooding, as well as many species of birds, mammals, and reptiles. It also helps buffer the coastline from storms. This indicator originates from LANDFIRE landcover.





Maritime forest

Not identified as maritime forest

Table 31: Indicator values for South Atlantic maritime forest within Virginia. A good condition threshold is not yet defined for this indicator.

	Indicator Values	Acres	Percent of Area
↑ High	Maritime forest	19,968	<0.1%
\downarrow Low	Not identified as maritime forest	3,084,482	11.3%
	Area not evaluated for this indicator	24,255,831	88.7%
	Total area	27,360,280	100%



This indicator uses remote sensing to calculate the unvegetated-vegetated ratio of tidal wetlands, which compares how much of a wetland is not covered by plants (e.g., sediment, rocks, open water) to how much is covered by plants. Marshes that maintain a higher proportion of vegetation tend to be more stable and resilient to threats like sea-level rise, erosion, and coastal development. This ratio, and how it changes over time, is a good surrogate for salt marsh degradation processes like sediment loss and conversion to open water. This indicator originates from a U.S. Geological Survey project on an unvegetated to vegetated ratio for coastal wetlands.





Stable coastal wetlands Other coastal wetlands

Not identified as coastal wetlands

Table 32: Indicator values for stable coastal wetlands within Virginia. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.

	Indicator Values	Acres	Percent of Area	
↑ High	Stable coastal wetlands	212,074	0.8%	
	Other coastal wetlands	136,657	0.5%	↑ In good condition
↓ Low	Not identified as coastal wetlands	931,667	3.4%	↓ Not in good condition
	Area not evaluated for this indicator	26,079,882	95.3%	
	Total area	27,360,280	100%	

Threats

Sea-level rise

NOAA's sea-level rise (SLR) inundation models represent areas likely to experience flooding at high tide based on each foot of SLR above current levels. Darker blue areas will experience flooding first, and at greater depth, compared to lighter blue areas. These models are not linked to a future timeframe; see the projections below. NOAA calculates the inundation footprint at "mean higher high water", or the average highest daily tide. The area covered in each SLR scenario includes areas projected to be inundated at lower levels. For example, the area inundated by 4 ft of SLR also includes areas inundated by 3 ft, 2 ft, 1 ft, and 0 ft of SLR (where 0 ft represents current levels).



To explore additional SLR information, please see NOAA's Sea Level Rise Viewer.



Flooding extent by projected sea-level rise (ft)



Table 33: Extent of flooding by projected average highest daily tide due to sea level rise within Virginia. Values from the <u>NOAA sea-level rise inundation data</u>.

Feet of sea-level rise	Acres	Percent of Area
0 feet	1,945,569	7.1%
1 foot	2,061,846	7.5%
2 feet	2,144,030	7.8%
3 feet	2,193,012	8.0%
4 feet	2,241,151	8.2%
5 feet	2,286,589	8.4%
6 feet	2,336,585	8.5%
7 feet	2,391,425	8.7%
8 feet	2,449,045	9.0%
9 feet	2,507,203	9.2%
10 feet	2,566,709	9.4%
Not projected to be inundated by up to 10 feet	5,392,743	19.7%
Sea-level rise unlikely to be a threat (inland counties)	19,400,731	70.9%
Sea-level rise data unavailable	97	<0.1%
Total area	27,360,280	100%

Table 34: Projected sea level rise by decade within Virginia. Values are based on area-weighted averages of decadal projections for 1-degree grid cells that overlap this area based on <u>NOAA's 2022 Sea Level Rise</u> <u>Report</u>. 2060 corresponds to the <u>SECAS goal</u>: a 10% or greater improvement in the health, function, and connectivity of Southeastern ecosystems by 2060.

SLR Scenario	2020 (ft)	2030 (ft)	2040 (ft)	2050 (ft)	2060 (ft)	2070 (ft)	2080 (ft)	2090 (ft)	2100 (ft)
Low	0.4	0.65	0.9	1.1	1.3	1.5	1.6	1.8	1.9
Intermediate- low	0.43	0.71	0.98	1.2	1.5	1.8	2	2.3	2.5
Intermediate	0.44	0.73	1	1.4	1.7	2.2	2.6	3.3	4
Intermediate- high	0.44	0.75	1.1	1.5	2.1	2.7	3.5	4.3	5.3
High	0.44	0.77	1.2	1.7	2.4	3.4	4.4	5.6	6.8

Urban growth

The FUTURES urban growth model predicts the likelihood that an area will urbanize at every decade from 2020 to 2100. Developed areas from the 2021 National Landcover Database serve as the baseline for current urban areas. The model simulates landscape change based on trends in population growth, local development suitability factors, and an urban patch-growing algorithm. It considers environmental drivers like distance to floodplain, slope, and available infrastructure, and even socio-economic status. The probability of urbanization for each area reflects how many times it urbanized out of 50 model runs.





Probability of urbanization by 2060

- Urban in 2021
- Very high likelihood of urbanization (>50% probability)
- High likelihood of urbanization (25 50% probability)
- Moderate likelihood of urbanization (2 25% probability)
- Not likely to urbanize

10.1% of this area is already urban in 2021, and an additional 11.0% has at least a moderate probability of urbanizing by 2060.

Table 35: Extent of projected urbanization by decade within Virginia. Values from <u>FUTURES model</u> projections for the contiguous United States developed by the <u>Center for Geospatial Analytics</u>, NC State University. 2060 corresponds to the <u>SECAS goal</u>: a 10% or greater improvement in the health, function, and connectivity of Southeastern ecosystems by 2060.

Decade	Acres	Percent of Area
Urban in 2021	2,771,595	10.1%
2030 projected extent	2,860,407	10.5%
2040 projected extent	2,906,823	10.6%
2050 projected extent	2,940,305	10.7%
2060 projected extent	2,972,486	10.9%
2070 projected extent	2,998,741	11.0%
2080 projected extent	3,017,621	11.0%
2090 projected extent	3,029,372	11.1%
2100 projected extent	3,034,869	11.1%
Not projected to urbanize by 2100	21,181,290	77.4%
Total area	27,360,280	100%

Ownership and Partners

Conserved lands ownership



Local

Ownership unknown

Designation

Table 36: Extent of ownership class within Virginia. Protected areas are derived from the <u>Protected Areas</u> <u>Database of the United States</u> (PAD-US v3.0) and include Fee, Designation, Easement, Marine, and Proclamation (Dept. of Defense lands only) boundaries. Note: areas are based on the polygon boundary of this area compared to protected area polygons, rather than pixel-level analyses used elsewhere in this report. Also note: PAD-US v3.0 includes protected areas that may overlap within a given area; this may cause the area within and between the following categories to be greater than the actual ground area.

Ownership	Acres	Percent of Area
Federal	2,135,298	7.8%
State/province	502,739	1.8%
Regional	15,620	<0.1%
Local	181,953	0.7%
Joint	388	<0.1%
Private non-profit conserved lands	315,602	1.2%
Private conservation land	1,151,796	4.2%
Tribal	1,959	<0.1%
Designation	1,207,518	4.4%
Ownership unknown	32,429	0.1%

Land protection status





- Managed for biodiversity (disturbance events proceed or are mimicked)
- Managed for biodiversity (disturbance events suppressed)
- Managed for multiple uses (subject to extractive uses such as mining or logging, or OHV use)
- No known mandate for biodiversity protection

Table 37: Extent of land protection status within Virginia. Protected areas are derived from the <u>Protected</u> <u>Areas Database of the United States</u> (PAD-US v3.0) and include Fee, Designation, Easement, Marine, and Proclamation (Dept. of Defense lands only) boundaries. Note: areas are based on the polygon boundary of this area compared to protected area polygons, rather than pixel-level analyses used elsewhere in this report. Also note: PAD-US v3.0 includes protected areas that may overlap within a given area; this may cause the area within and between the following categories to be greater than the actual ground area.

Land Protection Status	Acres	Percent of Area
Managed for biodiversity (disturbance events proceed or are mimicked)	277,299	1.0%
Managed for biodiversity (disturbance events suppressed)	970,215	3.5%
Managed for multiple uses (subject to extractive uses such as mining or logging, or OHV use)	2,750,308	10.1%
No known mandate for biodiversity protection	1,547,480	5.7%

Protected Areas

- George Washington and Jefferson National Forest (USDA FOREST SERVICE; 1,672,713 acres)
- SHEN (NPS; 195,789 acres)
- Mount Rogers (154,913 acres)
- Cumberland Forest (The Nature Conservancy; 153,444 acres)
- GREAT DISMAL SWAMP NATIONAL WILDLIFE REFUGE (Fee; 86,181 acres)
- Shenandoah Wilderness (82,336 acres)
- Fort A P Hill (74,612 acres)
- Marine Corps Base Quantico (58,924 acres)
- The Nature Conservancy Easement (PVT; 52,676 acres)
- NG Fort Pickett ARNG MTC (41,156 acres)
- Fort Pickett (VA Army National Guard; 41,156 acres)
- John H. Kerr Reservoir (37,318 acres)
- Virginia Department of Historic Resources Easement (PVT; 36,410 acres)
- BLRI (NPS; 35,237 acres)
- Goshen-Little North Mountain (VA Department of Wildlife Resources; 33,315 acres)
- Virginia Department of Forestry Easement (PVT; 28,561 acres)
- Little River (27,292 acres)

- Tidal Lands (Commonwealth of Virginia; 26,165 acres)
- Clinch Mountain (VA Department of Wildlife Resources; 24,854 acres)
- Bear Creek (23,396 acres)
- Fauquier County Easement (PVT; 21,229 acres)
- Assateague Island National Seashore (Unknown; 20,830 acres)
- ASIS (NPS; 20,362 acres)
- Appomattox-Buckingham State Forest (VA Dept of Forestry; 20,064 acres)
- APPA (NPS; 17,672 acres)
- ... and 7,555 more protected areas ...

Nearby land trusts

<u>Click here</u> to search for land trusts within 500 miles of this area on the Land Trust Alliance website.
Credits

This report was generated by the Southeast Conservation Blueprint Explorer, which was developed by <u>Astute Spruce, LLC</u> in partnership with the U.S. Fish and Wildlife Service under the <u>Southeast</u> <u>Conservation Adaptation Strategy</u>.

Data credits

Land ownership and conservation status is derived from the <u>Protected Areas Database of the United</u> <u>States</u> (PAD-US v3.0).

Future urban growth estimates derived from <u>FUTURES model projections for the contiguous United States</u> developed by the <u>Center for Geospatial Analytics</u>, NC State University.

Sea level rise data are derived from the National Oceanic and Atmospheric Administration's <u>Sea Level Rise</u> <u>Inundation Depth Data</u> and the <u>2022 Sea Level Rise Technical Report</u>.